



The Effects of High Mineral Alkaline Water Consumed over Three Consecutive Days on Reaction Time Following Anaerobic Exercise – A Randomized Placebo-Controlled Crossover Pilot Study

by

Michal Steffl¹, Ivana Kinkorova¹, Karolina Talar², Tereza Jandova¹,
Karolina Moulisova¹, Dan Omcirk¹, Jan Malecek¹, Jan Chrudimsky¹, Michał Wilk³,
Petr Stastny^{1,3}, Miroslav Petr^{1,3}

Drinking alkaline water after intense anaerobic exercise may enhance both cognitive and physical performance. This study aimed to investigate the effect of high mineral alkaline water consumed over three consecutive days on reaction time after anaerobic exercise in twelve healthy young males (aged 21.1 ± 1.3 years) with a valid sports medical examination. Participants were excluded when they took any medications or performance-enhancing supplements for the period of at least four weeks before the study commenced. Participants were randomly divided into two groups in this double-blind, placebo-controlled crossover pilot study. They ingested either alkaline water (AW) or regular table water (RTW) for three consecutive days before anaerobic exercise. The anaerobic exercise consisted of two 2-min high-intensity step-tests with a passive rest interval of 3 minutes between the two bouts of exercise. Performance in the step-test (W), reaction time for visual and auditory signals, the rate of perceived exertion (RPE), urine specific gravity, and lactate concentration were analysed. No effect of AW was found on reaction time and the other variables except anaerobic performance. Anaerobic performance was significantly higher after ingestion of AW in both step-tests ($p < 0.05$). The ingestion of AW for three consecutive days before anaerobic exercise seems to positively affect anaerobic performance.

Key words: Rate of Perceived Exertion, step-test, exercise, buffer system.

Introduction

Metabolic acidosis is a temporary consequence of an intense burst of exercise, such as a sprint by runners or cyclists, which limits an athlete's performance (Bangsbo et al., 1993; Robergs et al., 2004). The body's first line of defence against such change in internal pH is provided by buffer systems (protein, phosphate, and bicarbonate) when blood accepts hydrogen ions from the lactic acid produced in muscle tissue,

which is one of the major end products of anaerobic metabolism (Nelson and Cox, 2017). A shred of sufficient evidence indicates that alkaline agents can modify the blood buffering system by increasing the amount of bicarbonate concentration in the blood (increased blood pH), which in turn allows for a greater accumulation of lactate in the muscles during intense exercise and, thus leads to improved anaerobic performance

¹ - Faculty of Physical Education and Sport, Charles University, 16252 Prague, Czech Republic.

² - Faculty of Motor Rehabilitation, University of Physical Education in Krakow, 31-571 Krakow, Poland.

³ - The Jerzy Kukuczka Academy of Physical Education in Katowice, 40-065 Katowice, Poland.

(Sahlin, 1978; Zajac et al., 2009). For this reason, one of the recent interest in sports science has been the drinking of alkaline water (AW), which has been shown to enhance athletes' anaerobic performance, hydration status and increase both fasting and post-exercise arterial blood pH (Chycki et al., 2017, 2018; Kurylas et al., 2017; Ostojic and Stojanovic, 2014; Rocznioek et al., 2013). However, the exact alkalizing mechanism in terms of lactic acid accumulation and utilization after exercise is still unclear. Chycki et al. (2017, 2018) have shown that after drinking AW for 2-3 weeks, blood lactate levels at rest significantly decreased, but increased post-exercise, suggesting better lactate utilization.

Lactic acid has long been thought to be the culprit of muscle fatigue and pain as the anaerobic breakdown of glycogen leads to its intracellular accumulation. However, recent evidence on mammalian muscle has shown that reduced pH may have little influence on muscle contraction, and it is most likely the inorganic phosphate which is the major cause of muscular fatigue (Posterino et al., 2001; Westerblad, 2002, 2016). In the same way, there are currently contradictory opinions (Coco et al., 2009, 2016, 2020; Duncan et al., 2017; Mroczek et al., 2011; Tanaka et al., 2018; Tsukamoto et al., 2016) on how high blood lactate levels (induced either by exhaustive exercise or by an intravenous infusion of a lactate solution) may affect various psychomotor processes, especially simple reaction time.

Simple reaction time demands intense, simple attention and is considered an essential ability for many sport disciplines, especially the ones concerning visual stimulus (Cojocariu, 2011). For example, Perciavalle et al. (2014, 2016) have found that attentive capabilities and performance after an exhaustive exercise worsened, most likely due to increased blood lactate levels. Coco et al. (2020) also found increased reaction time and blood lactate after intense exercise, which were strongly correlated. On the other hand, Mroczek et al. (2011) found an initial decrease in reaction time despite increased blood lactate during and after volleyball sets.

Therefore, based on all the studies mentioned above, reporting both, improved anaerobic performance (linked to better lactate utilization and hydration status) after drinking AW and the effects of high blood lactate levels on reaction time, we investigated the effects of high

mineral AW consumed over three consecutive days on reaction time after anaerobic exercise, which is a novelty of this study.

Methods

Participants

Twelve young male students of the Faculty of Physical Education and Sport, Charles University, with an average age of 21.1 ± 1.3 , were recruited for the study. The inclusion criteria comprised increased anaerobic performance and cognition at a competitive level (e.g. a sub-elite competitive athlete), age between 18 – 25 years and current training of at least three times a week. All participants were classified as healthy and had a valid sports medical examination, which allowed them to participate in the experiment. They did not take any medications or any performance-enhancing supplements for at least four weeks before the study commenced (exclusion criterion). All participants signed informed consent after being informed about the study, including the possible risks. The study was conducted with the consent of the ethics committee of the Faculty of Physical Education and Sport, Charles University, Prague, Czech Republic, conforming to the ethical standards set by the Declaration of Helsinki (World Medical Association, 2013). Participants' characteristics are presented in Table 1.

Study Design

A pilot study with a double-blind, randomized, placebo-controlled crossover design was applied (Figure 1). Participants were randomly divided using free online statistical computing web programming (www.graphpad.com/quickcalcs) that generated the randomization schedule into two groups ($n = 6$) with different conditions: the consumption of 1) AW or 2) regular table water (RTW). At first all participants performed the baseline reaction time tests. A wash-out period of two weeks was applied to prevent a carry-over phenomenon before crossing-over. Participants were instructed to ingest 2.5 - 3 L (based on their body weight; i.e. at least $35 \text{ ml} \cdot \text{kg}^{-1}$ of body weight a day) of either AW or RTW (blinded) a day for three consecutive days (Kurylas et al., 2018). The water for the three-day consumption was provided in $6 \times 1.5 \text{ L}$ bottles; two bottles were to be consumed each day throughout the 24 hours (e.g. in the morning after awakening, during the day, and before going to

sleep). Before physical testing, lactate concentration was evaluated from capillary blood, and urine specific gravity was measured as well. The reaction time for visual and auditory signals was evaluated immediately after each physical test, along with the Rate of Perceived Exertion (RPE). The recovery time between the two physical tests was approximately 3 minutes, which corresponded to the execution time of the speed of reaction test. After finishing the second physical test, the urine analysis was performed again. The capillary blood was retaken in the fifth minute after the second step-test for evaluation of lactate concentration. To ensure similar conditions in terms of sports activity, sleeping time, and diet, participants were instructed to (1) omit strenuous physical activity the day before the examination; (2) three days before the testing itself, drink only the given amount of water (alkaline water or regular table water); (3) obey the requirement of 8 hours of sleep a day; (4) refrain from consuming alcohol, dietary supplements, and caffeinated products, and (5) follow an isocaloric mixed diet (55% carbohydrates, 20% protein, and 25% fat). Each trial was performed at the same time of the day to avoid circadian variations. The study lasted for a total of 3 weeks.

Alkaline Water

Commercially available glacial water with a pH of 8.8 was used in the study. The water was relatively highly alkaline compared to other commercially available products with high dissolved oxygen content ($\sim 8.5 \text{ mg}\cdot\text{l}^{-1}$) and low mineralization ($\sim 59 \text{ mg}\cdot\text{l}^{-1}$). It contained Na^+ $9.52 \text{ mg}\cdot\text{l}^{-1}$; Ca^{2+} $4.47 \text{ mg}\cdot\text{l}^{-1}$; Mg^{2+} $1.66 \text{ mg}\cdot\text{l}^{-1}$; Al^+ $0.0193 \text{ mg}\cdot\text{l}^{-1}$, and Cl^- $8.18 \text{ mg}\cdot\text{l}^{-1}$, other ions were present in small amounts. The total hardness of water was $0.180 \text{ mmol}\cdot\text{l}^{-1}$. Odour and taste were indistinguishable from the RTW.

Physical Test

A modified version of the high-intensity step-test was used in the study (Perciavalle et al., 2016). In this study, we used a 50 cm high step, and we set the execution time to 2 minutes. Ascents and descents were performed alternately with the right and the left leg, where one leg always remained on the podium, and the tested person was not allowed to rest hands on thighs. An authorized member of the research staff supervised the correct technical execution of the test, tracked the time, motivated the participants, and recorded the results. The

performance in watts (W) was calculated as follow:

$$P = \frac{F \times s}{t} = \frac{m \times 9.81 \times h \times c \times 1.33}{t}$$

where t = test duration [s]; F = force [N]; s = distance [m]; m = body weight [kg]; h = step height [m]; c = number of steps; 1.33 = coefficient for positive (1.0) and negative work (0.33).

Reaction Time

Special software developed for this purpose at the Department of Sports Diagnostics, at the Faculty of Physical Education and Sport Charles University, was used to measure reaction time. Briefly, the tested person had to press a button connected to the computer when a visual (on the screen of the computer) or an auditory signal (beep) appeared. The tested person had to first react as soon as possible to twenty visual and then to twenty acoustic signals for 80 s. The five fastest and five slowest reaction times were automatically deleted from the average reaction time calculation. To avoid settling habituation, signals were randomized with intervals spanning between 1 and 3 s. The reaction time in [s] was measured for visual and auditory signals separately.

Blood and Urine Tests

20 μl of capillary blood was collected from the fingertip to determine lactate concentration before and after exercise. Blood samples were analysed by the analyser SUPER GL - Biovendor. Urine specific gravity before and after exercise was analysed using the TIPA LH-T50 Refractometer.

Rate of Perceived Exertion

The 10 point Rate of Perceived Exertion (RPE) scale was used in this study (Borg, 1982). The RPE is a self-reported scale ranging from 1 = extremely easy to 10 = extremely hard exercise, and the participants were familiarized with the scale before the start of the experiment.

Data Analysis

Our internal reliability data measured by the intra-class correlation coefficient (ICC) for both exercise and reaction time protocols indicated an excellent level of reliability for all the considered variables (ICC ranging from 0.96-0.99). Means and standard deviations (SDs) were calculated for each variable. Because of the small sample size, we used non-parametric statistical methods for repeated measures (Sullivan et al., 2016; Yan et al., 2017). The Friedman test ($\alpha = 0.05$) with the Wilcoxon signed

ranks test with Bonferroni correction in the post-hoc analysis were used to test our hypotheses. In the Bonferroni correction, the standard level of significance was divided by the number of comparisons $\alpha = 0.05/6 = 0.008$; therefore, the statistically significant result after the correction was when $p < 0.008$. All the statistics were carried out in the statistical software IBM SPSS Statistics 24. Graphs for the reaction time were created separately in MS Excel.

Results

There were no significant changes observed in reaction time for both visual and auditory signals after drinking AW or RTW when compared to the resting values. Graphical representation of both reaction times is provided in Figure 2.

Performance measured in watts (W) in the first (853.2 ± 128.7 vs. 739.6 ± 107.6) and second (814.4 ± 127.6 vs. 688.6 ± 86.1) step-tests was

significantly higher after the intake of AW when compared to RTW, respectively. Performance in the second step-test was not significantly different from the first step-test after both AW and RTW.

There was no significant difference in the RPE between both groups after both tests. The RPE significantly increased ($p = 0.004$) after the second step-test (when considering the first step-test values) in RTW. There was a significant difference between the first measurement in AW and the second measurement in RTW.

No significant difference was observed between the initial lactate concentration after AW and after RTW. There was a significant increase in lactate concentration after the step-tests under both conditions.

No significant difference was observed in urine specific gravity.

Graphical representation of performance in step-tests, RPE, lactate concentration and urine specific gravity is provided in Figure 3.

Table 1.
Descriptive statistics of the study participants presented as means and standard deviations.

| | N = 12 |
|--------------------------|-------------|
| Age (years) | 21.1 (1.3) |
| Body height (cm) | 180.8 (5.3) |
| Body mas (kg) | 76.4 (4.9) |
| BMI (kg/m ²) | 23.3 (1.0) |




Figure 1.

The course of the study.

Note. AW - alkaline water; RTW - regular table water.




Figure 2. Reaction time changes to visual $p = 0.685$ (and auditory signals) $p = 0.249$ after ingesting alkaline water.
 Note. AW - alkaline water; RTW - regular table water.




Figure 3 The effect of AW on performance in the step-test, RPE, lactate concentration, and urine specific gravity.
 Note. AW - alkaline water; RTW - regular table water.

Discussion

In this study, we investigated the effects of high mineral alkaline water consumed over three consecutive days on reaction time after anaerobic exercise. This study found no effect on reaction time for both visual and auditory signals in healthy young males. However, a significant difference in anaerobic performance (measured by a step-test) was observed.

Recent studies have found a positive effect of alkaline water ingestion on anaerobic performance, which stems from improving the acid-base balance, lactate utilization and hydration (Chycki et al., 2017, 2018; Kurylas et al., 2017). We, therefore, hypothesized that it might also affect attentive performance, such as simple reaction time, since several recent studies have linked high blood lactate levels to various psychomotor processes both positively and negatively (Coco et al., 2009, 2016, 2020; Mroczek et al., 2011; Perciavalle et al., 2014).

In our study, reaction time did not change when compared to pre-exercise values, which can be considered positive, since athletes with high levels of blood lactate even at rest, had attentional performances, including reaction time-limited (Coco et al., 2009). In terms of the reaction time and lactate concentration, the results of our study partially support the findings of Mroczek et al. (2011) who measured reaction time at rest and blood lactate concentration pre-game and during four volleyball sets and found that reaction time decreased during the first set significantly when compared to the pre-game test and then remained at the same level at all time. In their study, the association between reaction time and lactate was not calculated, due to the study's nature. However, future and more extensive studies conducted preferably on different populations are needed to investigate this phenomenon further.

Generally, our study results are more likely to be attributed to the good hydration status in both groups. Good hydration can have a significant positive influence on aerobic and anaerobic performance, including people's mental state. It has been shown that even a minor 1% hypohydration influences a variety of cognitive functions (Karpęcka and Frączek, 2020), specifically the speed of reaction (Cian et al., 2001; Tomporowski et al., 2007). Indeed, reaction time is controlled by the central nervous system and is

affected by many factors, including age, gender, stimulus (type and intensity), arousal, fatigue and exercise (Cote et al., 2009; Der and Deary, 2006; Kashihara and Nakahara, 2005). For example, Tsukamoto et al. (2016) showed an improved executive function, including reaction time after high-to-moderate intensity interval exercise, although Perciavalle et al. (2014, 2016) and Coco et al. (2016; 2020) found the opposite. Nevertheless, in our study, we intended to investigate only the differences in reaction time after the three-day ingestion of alkaline water vs. tap water to provide pilot data to support future studies.

In our study, we found significantly higher performance in the two step-tests after the three-day ingestion of alkaline water. In terms of lactate concentration (measured by capillary blood), our initial levels did not differ significantly between the two groups; even though an increased lactate level is noted following the ingestion of alkaline water, which is in partial agreement with Chycki et al. (2018) and Kurylas et al. (2018). The muscle pH of trained athletes can drop from a resting value of 7.0 down to 6.4 following intense exercise, and bicarbonate-CO₂ accounts for more than 90% of the plasma buffering capacity (Robergs et al., 2004). In our study, we did not find any significant increments in urine specific gravity, unlike in the two previous studies (Chycki et al., 2017; Kurylas et al., 2018), which was something expected (three consecutive days vs. six consecutive weeks (Kurylas et al., 2018)). Our results have instead served to check the hydration status of both groups, which was nearly identical, suggesting good participant's compliance, and as mentioned above, it may be the main reason why reaction time was unchanged in both groups.

Nevertheless, the six-week study (Kurylas et al., 2018) showed that highly alkalized water delayed muscular acidosis during anaerobic exercise, acted as prevention of dehydration, and improved recovery. As the RPE scale can 'indirectly' assess all physiological signs (the sensation of fatigue and pain after such exercise) of redox homeostatic mechanisms (Borg, 1982), it appears that this greater muscle buffering capacity and enhanced removal of protons, resulting in increased glycolytic ATP production, may be responsible for the improved anaerobic performance following alkaline water ingestion. Such an effect of alkaline water ingestion before

anaerobic exercise is similar to sodium or potassium bicarbonate supplementation. The use of sodium bicarbonate as an alkaline agent can be dated back to 1928 when Haldane and his physiologist colleague, Davies (Haldane, 1928), decided to experiment on themselves how the body controls blood pH. Since then, the ingestion of sodium bicarbonate has proven effective in speed endurance sports; however, its use has been limited as it often causes gastrointestinal distress or metabolic alkalosis due to sodium overload (McNaughton et al., 2008). These supplements have the potential to modify the adaptive response to endurance training by affecting acid-base balance, redox status, reactive oxygen species (ROS) signalling, or cumulative training load, thereby affecting the cellular signalling responses to training (Rothschild and Bishop, 2020). Therefore, the idea of increasing the capacity of the body's buffering system by ingesting alkaline agents to enhance anaerobic performance stems from the ionic biochemical interplay (Chycki et al., 2018; Coco et al., 2016). Although there is a large body of evidence to support the use of sodium bicarbonate for sports performance, its use has been associated with possible gastrointestinal side effects (Kahle et al., 2013). Contrary to this, it seems that alkaline water can serve as an effective alternative to sodium bicarbonate without the mentioned side effects, improving an athlete's anaerobic performance (Chycki et al., 2017, 2018; Kurylas et al., 2018; Ostojic and Stojanovic, 2014). Future studies could consider comparing the use of sodium

bicarbonate, alkaline water, as well as an isotonic beverage to assess not only the negative side effects, but also anaerobic performance following these three supplements.

It should be noted that some methodological considerations are involved in this study. First, the main limitation of this study is the small sample size due to the pilot nature of the investigation. For this reason, we used non-parametric statistical testing as the recommended approach in medical and biomedical research (Sullivan et al., 2016; Yan et al., 2017), along with the calculation of effect sizes to account for this problem. Another limitation is that this study did not measure the acid-base balance, which, otherwise, could provide a more detailed mechanistic insight into our findings. Future studies should ideally focus on measuring this variable alongside and comparison to other food supplements like hydrogen-rich water (Timón et al., 2021) or bicarbonates (Chycki et al., 2021). In our study, we used natural AW with pH 8.8, which is not the same as extremely high AW with pH 10, for which negative health consequences in rats have been reported (Merne et al., 2001; Watanabe et al., 1997).

Conclusions

In conclusion, the results of the present study indicate that ingestion of high mineral alkaline water for short, three-day duration has no effect on reaction time. However, alkaline water seems to positively affect anaerobic performance, which needs to be further confirmed by future studies with a larger number of subjects.

Acknowledgements

This research was supported by research grants from the Charles University, Czech Republic (UNCE/HUM/032 and Q41). Student funding was provided by internal university funding SVV 2020-2022-260599.

References

- Bangsbo J, Johansen L, Graham T, Saltin B. Lactate and H⁺ effluxes from human skeletal muscles during intense, dynamic exercise. *J Physiol*, 1993; 462: 115-133
- Borg GA. Psychophysical bases of perceived exertion. *Med Sci Sports Exerc*, 1982; 14: 377-381
- Chycki J, Kurylas A, Maszczyk A, Golas A, Zajac A. Alkaline water improves exercise-induced metabolic acidosis and enhances anaerobic exercise performance in combat sport athletes. *PLoS One*, 2018; 13: e0205708
- Chycki J, Zajac T, Maszczyk A, Kurylas A. The effect of mineral-based alkaline water on hydration status and the metabolic response to short-term anaerobic exercise. *Biol Sport*, 2017; 34: 255-261

- Chycki J, Zajac A, Toborek M. Bicarbonate supplementation via lactate efflux improves anaerobic and cognitive performance in elite combat sport athletes". *Biol Sport*. 2021; 38(4): 545–553
- Cian C, Barraud PA, Melin B, Raphel C. Effects of fluid ingestion on cognitive function after heat stress or exercise-induced dehydration. *Int J Psychophysiol*, 2001; 42: 243-251
- Coco M, Buscemi A, Guerrera CS, Di Corrado D, Cavallari P, Zappala A, Di Nuovo S, Parenti R, Maci T, Razza G, Petralia MC, Perciavalle V, Perciavalle V. Effects of a bout of intense exercise on some executive functions. *Int J Environ Res Public Health*, 2020; 17: 898
- Coco M, Di Corrado D, Calogero RA, Perciavalle V, Maci T, Perciavalle V. Attentional processes and blood lactate levels. *Brain Res*, 2009; 1302: 205-211
- Coco M, Perciavalle V, Cavallari P, Perciavalle V. Effects of an exhaustive exercise on motor skill learning and on the excitability of primary motor cortex and supplementary motor area. *Medicine (Baltimore)*, 2016; 95: e2978
- Cojocariu A. Measurement of reaction time in Qwan Ki Do. *Biol Sport*, 2011; 28: 139-143.
- Cote KA, Milner CE, Smith BA, Aubin AJ, Greason TA, Cuthbert BP, Wiebe S, Duffus SE. CNS arousal and neurobehavioral performance in a short-term sleep restriction paradigm. *J Sleep Res*, 2009; 18: 291-303
- Der G, Deary IJ. Age and sex differences in reaction time in adulthood: results from the United Kingdom Health and Lifestyle Survey. *Psychol Aging*, 2006; 21: 62-73
- Duncan MJ, Clarke ND, Cox M, Smith M. The influence of cycling intensity upon cognitive response during inferred practice and competition conditions. *J Sports Sci*, 2017; 35: 1865-1871
- Haldane JBS. *Possible worlds and other papers*. New York, London: Harper & Brothers Publishers; 1928
- Kahle LE, Kelly PV, Eliot KA, Weiss EP. Acute sodium bicarbonate loading has negligible effects on resting and exercise blood pressure but causes gastrointestinal distress. *Nutr Res*, 2013; 33: 479-86.
- Kashihara K, Nakahara Y. Short-term effect of physical exercise at lactate threshold on choice reaction time. *Percept Mot Skills*, 2005; 100: 275-291
- Karpecka E, Frączek B. Macronutrients and water – do they matter in the context of cognitive performance in athletes?. *Balt J Health Phys Activ*, 2020; 12: 114-124
- Kurylas A, Zajac T, Chycki J, Maszczyk A. Anaerobic performance and acid-base balance in basketball players after the consumption of highly alkaline water. *Int J Food Sci Nutr*, 2018; 5: 134 - 139
- Kurylas A, Zajac T, Zydek G. The Effectiveness of alkaline water in hydrating athletes. *J Nutrition Health Food Sci*, 2017; 5: 1 - 4
- McNaughton LR, Siegler J, Midgley A. Ergogenic effects of sodium bicarbonate. *Curr Sports Med Rep*, 2008; 7: 230-236
- Merne ME, Syrjanen KJ, Syrjanen SM. Systemic and local effects of long-term exposure to alkaline drinking water in rats. *Int J Exp Pathol*, 2001; 82: 213-219
- Mroczek D, Kawczynski A, Chmura J. Changes of reaction time and blood lactate concentration of elite volleyball players during a game. *J Hum Kinet*, 2011; 28: 73-78
- Nelson DL, Cox MM. *Loose-leaf version for Lininger principles of biochemistry*. London: W. H. Freeman; 2017
- Ostojic SM, Stojanovic MD. Hydrogen-rich water affected blood alkalinity in physically active men. *Res Sports Med*, 2014; 22: 49-60
- Perciavalle V, Di Corrado D, Scuto C, Perciavalle V, Coco M. Attention and blood lactate levels in equestrians performing show jumping. *Percept Mot Skills*, 2014; 118: 733-745
- Perciavalle V, Marchetta NS, Giustiniani S, Borbone C, Perciavalle V, Petralia MC, Buscemi A, Coco M. Attentive processes, blood lactate and CrossFit((R)). *Phys Sportsmed*, 2016; 44: 403-406
- Posterino GS, Dutka TL, Lamb GD. L(+)-lactate does not affect twitch and tetanic responses in mechanically skinned mammalian muscle fibres. *Pflugers Arch*, 2011; 442: 197-203
- Robergs RA, Ghiasvand F, Parker D. Biochemistry of exercise-induced metabolic acidosis. *Am J Physiol Regul Integr Comp Physiol*, 2004; 287: R502-516
- Roczniok R, Maszczyk A, Stanula, Stanula A, Czuba M, Pietraszewski P, Kantyka J Starzynski M. Physiological and physical profiles and on-ice performance approach to predict talent in male youth ice hockey players during draft to hockey team. *Isokinet Exerc Sci*, 2013; 21(2):121-127
- Rothschild JA, Bishop DJ. Effects of dietary supplements on adaptations to endurance training. *Sports Med*, 2020; 50: 25-53

- Sahlin K. Intracellular pH and energy metabolism in skeletal muscle of man. With special reference to exercise. *Acta Physiol Scand Suppl*, 1978; 455: 1-56
- Sullivan LM, Weinberg J, Keaney JF, Jr. Common statistical pitfalls in basic science research. *J Am Heart Assoc*, 2016; 5
- Tanaka Y, Saihara Y, Izumotani K, Nakamura H. Daily ingestion of alkaline electrolyzed water containing hydrogen influences human health, including gastrointestinal symptoms. *Med Gas Res*, 2018; 8: 160-166
- Timón R, Olcina G, González-Custodio A, Camacho-Cardenosa M, Camacho-Cardenosa A, Guardado I. Effects of 7-day intake of hydrogen-rich water on physical performance of trained and untrained subjects. *Biol Sport*, 2021; 38: 269-275. doi:10.5114/biolsport.2020.98625
- Tomprowski PD, Beasman K, Ganio MS, Cureton K. Effects of dehydration and fluid ingestion on cognition. *Int J Sports Med*, 2007; 28: 891-896
- Tsukamoto H, Suga T, Takenaka S, Tanaka D, Takeuchi T, Hamaoka T, Isaka T, Hashimoto T. Greater impact of acute high-intensity interval exercise on post-exercise executive function compared to moderate-intensity continuous exercise. *Physiol Behav*, 2016; 155: 224-230
- Watanabe T, Kishikawa Y, Shirai W. Influence of alkaline ionized water on rat erythrocyte hexokinase activity and myocardium. *J Toxicol Sci*, 1997; 22: 141-152
- Westerblad H. Acidosis Is Not a Significant Cause of Skeletal Muscle Fatigue. *Med Sci Sports Exerc*, 2016; 48: 2339-2342
- Westerblad H, Allen DG, Lannergren J. Muscle fatigue: lactic acid or inorganic phosphate the major cause? *News Physiol Sci*, 2002; 17: 17-21
- World Medical Association. World Medical Association Declaration of Helsinki: ethical principles for medical research involving human subjects. *JAMA*, 2013; 310(20): 2191-2194
- Yan F, Robert M, Li Y. Statistical methods and common problems in medical or biomedical science research. *Int J Physiol Pathophysiol Pharmacol*, 2017; 9: 157-163
- Zajac A, Cholewa J, Poprzecki S, Waskiewicz Z, Langfort J. Effects of sodium bicarbonate ingestion on swim performance in youth athletes. *J Sports Sci Med*, 2009; 8: 45-50

Corresponding author:**Petr Stastny,**

Faculty of Physical Education and Sport, Charles University, 16252 Prague, Czech Republic
The Jerzy Kukuczka Academy of Physical Education in Katowice, 40-065 Katowice, Poland
E-mail: stastny@ftvs.cuni.cz