

High-Intensity Interval Training and Moderate-Intensity Continuous Training Affect Running Economy in Endurance Runners: A Systematic Review and Meta-Analysis of Randomized Controlled Trials

by

Yangya Feng ^{1,2}, Daxin Li ¹, Yangli Liu ¹, Donghui Tang ^{1,*}

This systematic review evaluated the effects of high-intensity interval training (HIIT) and moderate-intensity continuous training (MICT) on running economy (RE) in endurance runners. The search was completed in March 2024 based on research databases, and the language of publication was restricted to English. The primary outcome measure was RE, and it was categorized into three subgroups: Zone 1 (Z1), Zone 2 (Z2), and Zone 3 (Z3). The secondary outcomes assessed were maximal oxygen uptake ($VO_{2\max}$) and blood lactate concentration. HIIT significantly improved RE compared to MICT ($SMD = 0.44$, 95% CI [0.15, 0.72], $Z = 3.01$, $p < 0.05$). MICT showed a greater effect on $VO_{2\max}$ ($MD = 2.48$, 95% CI [1.61, 3.34], $Z = 5.60$, $p < 0.05$). HIIT was more effective at reducing blood lactate levels ($MD = -0.15$, 95% CI [-0.28, -0.02], $Z = 2.20$, $p < 0.05$). The results indicate that HIIT was more effective than MICT in enhancing RE and delaying lactate accumulation. HIIT can further improve RE and postpone blood lactate accumulation when performed at or below the lactate threshold ($\leq Z2$). $VO_{2\max}$ was more pronounced with MICT. These findings suggest that endurance runners and coaches should choose appropriate methods to optimize physiological adaptations.

Keywords: athletes; endurance; training methods; aerobic metabolism

Introduction

In endurance running, the efficiency and economy of energy utilization in human movement are crucial factors that impact performance. This efficiency can be quantified through running economy (RE). RE is defined as the energy expenditure associated with running at submaximal intensities (specific running speeds) and the ratio of work output to oxygen consumption (Saunders et al., 2004; Shaw et al., 2013). It is typically measured by oxygen uptake per unit of body weight over time during exercise ($O_2 \text{ ml} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$) or by oxygen uptake per unit of body weight per distance ($O_2 \text{ ml} \cdot \text{kg}^{-1} \cdot \text{km}^{-1}$) (Jones et al., 2021).

Traditional measures of RE have typically been assessed in laboratory settings by having

athletes run on a treadmill. In recent years, portable oxygen analyzers have gained widespread use in outdoor settings, providing significant convenience for athletes. RE is influenced by various factors, including the environment, sports biomechanics, physiology, and others (Saunders et al., 2004; Van Hooren et al., 2024). The spring-mass model, in which the spring-like action of the supporting leg counteracts the body's bounce upon contact with the ground, is a crucial element of RE. Mechanical energy is stored in the muscles, tendons, and ligaments that function across joints. Runners with high running economy (RE) utilize less oxygen at the same steady-state speed compared to runners with low RE (Black et al., 2018). Numerous studies have indicated that trained individuals are more

¹ College of P.E. and Sport, Beijing Normal University, Beijing, China.

² College of P.E. and Sport, Qinghai Normal University, Xining, China.

* Correspondence: b2022310700@126.com

economical than their untrained or less-trained counterparts (Morgan et al., 1995; Unhjem, 2024). It has been demonstrated that training enhances both the morphology and functionality of skeletal muscle mitochondria. At a given submaximal running speed, trained runners utilize less oxygen per mitochondrial respiratory chain due to enhanced respiratory capacity in skeletal muscle. These adaptations lead to a slower rate of muscle glycogen consumption in the active musculature, reduced disruption of homeostasis, and improvements in RE (Holloszy et al., 1977).

In long-distance running, many athletes utilize high-intensity interval training (HIIT) and moderate-intensity continuous training (MICT) to enhance their aerobic capacity and overall athletic performance. HIIT can activate larger motor units, increase the recruitment of fast oxidative and glycolytic muscle fibers (Jastrzębski et al., 2025; Lambole et al., 2020; Laursen et al., 2007;), induce lactic acid production to stimulate both muscles and the cardiovascular system, enhance muscle pumping function and stroke volume (Dolan et al., 2024; Rakobowchuk et al., 2008; Ramos et al., 2015; Seo et al., 2024), increase mitochondrial volume, and reduce plasma H⁺ concentration (Perry et al., 2008; Ruegsegger et al., 2023). MICT recruits a greater proportion of slow-twitch muscle fibers, enhances mitochondrial content, and increases the oxidative capacity of muscle fibers (Skelly et al., 2021). It also boosts muscle capillary density and improves myocardial perfusion and oxygen delivery (Cocks et al., 2016). HIIT can activate signaling pathways such as peroxisome proliferator-activated receptor gamma coactivator 1-alpha (PGC-1 α), a key regulator of mitochondrial biogenesis (Torma et al., 2019), and adenosine monophosphate (AMP) activated kinase (AMPK), an important energy-sensing molecule (Gibala, 2009; Kjøbsted et al., 2018). HIIT promotes mitochondrial biogenesis (Dohlmann et al., 2018) and induces various microRNAs, including miR-486-5p, miR-208b, and miR199a-3p, which significantly enhance muscle protein synthesis and hypertrophy (D'Souza et al., 2018; Grieb et al., 2023). Additionally, HIIT improves fat oxidation and oxygen uptake (Atakan et al., 2022). MICT can activate PGC-1 α through calcium/calmodulin-dependent protein kinase II (CaMKII) and the phosphatase calcineurin A (CnA), thereby regulating various downstream targets (Furrer et

al., 2023; Laursen, 2010). MICT has a more pronounced effect on the expression of miRNAs, including miR-21, miR-133, miR-19b-3p, and miR-503-5p (Grieb et al., 2023; Widmann et al., 2022). These microRNAs (miRNAs) can regulate myoblast proliferation and differentiation, enhance the oxidative metabolism of slow-twitch muscle fibers, and promote muscle adaptation (Gonçalves et al., 2019; Grieb et al., 2023; Massart et al., 2021; Widmann et al., 2022). Both training modes induce complex changes in microRNAs, with some being regulated exclusively by MICT or HIIT (Grieb et al., 2023; Widmann et al., 2022). HIIT is frequently utilized as an alternative to MICT (Cocks et al., 2016). Consequently, HIIT and MICT have emerged as the most prevalent endurance training strategies (González-Mohíno et al., 2020).

While there is scientific evidence demonstrating that endurance training reduces oxygen uptake, whether interval training is more effective than continuous training remains a debated issue in the literature (Cocks et al., 2016; González-Mohíno et al., 2020). MICT can enhance the body's capacity to transport and utilize oxygen, improve cardiovascular efficiency, and decrease oxygen uptake during exercise by increasing the adaptability of physiological factors, such as mitochondrial content and capillary density (Cocks et al., 2016; Hesketh et al., 2019). HIIT can induce changes in various physiological factors, including lactic acid accumulation, glycolysis, and oxidative utilization, through varying interval intensities and continuous training protocols (Morais et al., 2025; Sperlich et al., 2010). These changes, in turn, affect mitochondrial function and related proteases, ultimately influencing running economy (González-Mohíno et al., 2020; Schaun et al., 2018). There is no consensus regarding the effects of HIIT and MICT on running economy (González-Mohíno et al., 2020; Wang and Wang, 2024). Several studies have investigated the effects of HIIT and MICT on improving running economy (Schaun et al., 2018; Skovgaard et al., 2018a). Some studies employing similar training strategies found no significant improvements (Parmar et al., 2021; Wang and Wang, 2024). Other researchers have noted that improvements in running economy occur only with low-intensity training (González-Mohíno et al., 2020; Zinner et al., 2018). It remains uncertain whether HIIT has a comparable impact on running economy. This

study investigated the effects of HIIT and MICT, summarizing the changes in physiological indices such as maximal oxygen uptake, running economy, and lactate threshold to provide insights into scientific training and heretical basis for endurance runners.

Methods

Search Strategy

The processes outlined here were based on the PRISMA Statement (Page et al., 2021), and the registration number is INPLASY2024110120. We searched electronic databases, including PubMed, Embase, Scopus, and Web of Science, from their inception to March 2024 using the following search phrases along with Boolean operators: (running OR jogging OR "marathon running" OR "distance running" OR "endurance running" OR "distance runners" OR "endurance runners" OR "middle distance runners") AND ("running economy" OR "energy metabolism" OR "metabolism energy" OR "energy expenditure" OR "energy expenditures" OR "expenditure energy" OR energy OR "energy cost" OR "metabolic cost") AND ("intermittent training" OR "interval exercise" OR "interval running" OR "sprint interval training" OR "intensity training"). The language of publication was restricted to English, and the publication type was restricted to randomized controlled trials (RCTs).

Inclusion and Study Selection

Inclusion criteria were developed according to PICOS principles. Studies that satisfied the following criteria: (1) participants were endurance runners (P), (2) the intervention was interval training (I) compared with continuous training (C) in the control group, (3) studies reported outcomes related to running economy or energy consumption (O), (4) RCTs published in journals with peer review (S), (5) the study utilized a run-based test course or a run-based training program; were considered in the present review and meta-analysis.

Studies were excluded if they (1) lacked a specified intervention type or were not randomized controlled trials, (2) did not adhere to the established guidelines for a training design, (3) involved training interventions lasting less than four weeks (this criterion was implemented because neuromuscular adaptations have been

observed in trained individuals within as little as four weeks, the surface electromyography activity of muscles and the hypertrophy of muscles increased significantly at the 4th week) (Baroni et al., 2013; Del Vecchio et al., 2019; Mayhew et al., 1995; Moritani and deVries, 1979), or if they involved non-interval training interventions. Furthermore, this systematic review did not include any review articles.

Outcome Measurements

The test results included running economy (RE) as the primary outcome, which was calculated using energy cost or steady-state oxygen consumption. Secondary outcome measures included at least one of the following: VO_{2max} and/or blood lactic acid levels.

Data Collection Process

The included studies were screened by the Endnote X9 citation manager (Liu et al., 2024). Two researchers independently reviewed this literature, excluded duplicate and irrelevant studies, and then independently decided whether to include them in the study. In the event of disagreement, a third researcher was consulted to make the final decision. First, the titles and abstracts were reviewed to exclude studies that clearly did not meet the inclusion criteria. Then, the full texts of the remaining manuscripts were screened and read, the eligibility was evaluated, and the final literature to be included was determined. Two researchers independently extracted studies that met the inclusion and exclusion criteria, including article publication information, participants' characteristics, the study design, the research content, intervention-related information, and RE results. The authors of the considered papers were contacted for clarification of incomplete or missing data.

We used Excel to store extracted data. Detailed data were registered, including the first author, the publication year, the country of publication, subjects' characteristics, VO_{2max}, training interventions, methods, running economy, and other relevant factors. We utilized the post-intervention mean and standard deviation comparison between groups for calculating effect sizes. When no standard deviations (SDs) were available, they were calculated from standard errors (SEs), CIs, t or p values.

Risk of Bias Assessment

Two reviewers independently assessed the risk of bias, and any disagreements were resolved through discussion with a third author. The bias in the included randomized controlled trials was evaluated using the Cochrane Risk of Bias Tool, which categorized each item as of low risk, high risk, or unclear risk of bias concerning random sequence generation, allocation concealment, blinding of participants and personnel, blinding of outcome assessment, incomplete outcome data, selective reporting, and other biases (Higgins and Green, 2011).

The performance bias item was eliminated because it was not practicable to blind the participants. A technique was established to assess the potential for bias in selection (random sequence generation and allocation concealment), detection, attrition, and reporting biases. The study was classified as of "high risk" or "low risk", and if the information was omitted, it was graded as "unclear". We addressed the impact of bias levels on the weights in our analyses by conducting sensitivity analyses that excluded studies with high risk of bias, and we found that the overall results remained robust. When a significant result ($p < 0.05$) was observed, a funnel plot test was also performed to assess publication bias. The effects of both HIIT and MICT on running economy performance were studied through the funnel plot.

Statistical Analyses

We conducted a meta-analysis using RevMan 5.3.5 to determine whether HIIT affected running economy performance. For continuous outcomes, we utilized the sample size, mean, and standard deviation (SD) values from both the experimental and control groups. We calculated Hedges' g , which represented standardized mean differences (SMD), to assess the magnitude of changes in outcomes between the pre-training and post-training periods (Nakagawa and Cuthill, 2007). Additionally, we employed inverse variance weighting to assign weights to each study included in the meta-analysis, with studies contributing more to the pooled effect size when they were of higher quality (e.g., larger sample size, lower risk of bias). To minimize the influence of studies with high risk of bias, lower weights were assigned to studies with methodological limitations or small sample sizes. A random effects model was utilized

in the forest plot to synthesize the effect sizes for each group. We aggregated these effect sizes according to the criteria outlined in the Cochrane Handbook when the included publications assessed running economy at various velocities or included multiple training groups (Higgins et al., 2023). Then, statistical heterogeneity was discussed using the I^2 statistic, where values of 25%, 50%, and 75% indicated low, moderate, and high heterogeneity, respectively (Higgins et al., 2003). The synthesis of the included trials was considered to exhibit significant heterogeneity if $p < 0.1$ or $I^2 > 0.5$, prompting the use of a random effects model. To further assess the robustness of the findings and ensure that potential bias did not significantly affect the pooled results, sensitivity analyses were conducted by systematically excluding each trial one by one. The results were visualized using a funnel plot, which appeared symmetric, suggesting no significant publication bias. Additionally, subgroup analyses were performed to explore potential sources of heterogeneity based on study characteristics, including the sample size, training intensity, and study quality. This process helped to determine whether the overall findings were influenced by any particular subset of studies with higher risk of bias.

Results

Study Selection

A total of 1,421 articles were retrieved from four databases: PubMed, Embase, Web of Science, and Scopus. After removing duplicates using EndNote software, 673 articles remained. Following further screening of titles and abstracts, 57 full-text publications were retained for detailed review. Of these studies, 12 lacked controlled outcomes, 15 did not address running economy (RE) or energy expenditure, 11 focused on intervention-related injuries, 6 involved mixed training interventions, and 4 had training duration of less than four weeks. These articles were excluded, leaving 9 studies that were included in the systematic review.

Study Characteristics

Tables 1 and 2 summarize the studies and outline the key characteristics of the participants involved. The nine studies that met the inclusion criteria are detailed, including their participants' profiles and running economy data. A total of 168

participants (110 males and 48 females) were included in the analysis, with age ranging from 21.9 to 46.9 years. Maximal oxygen uptake data were recorded, with values ranging from 34.0 to 71.4 $\text{ml}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$. Based on the weighted mean values from the studies reporting participants' characteristics for each group, minimal differences at baseline were observed for age (38.2 vs. 39.3 years), body mass (81.6 vs. 82.2 kg), body height (1.82 vs. 1.81 m), and $\text{VO}_{2\text{max}}$ (59.6 vs. 59.3 $\text{mL}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$) between the control and HIIT groups. Seven studies included moderately trained or recreational runners, one study featured well-trained participants, and another examined highly trained runners. Participants were involved in a variety of activities, including triathlons and endurance or recreational distance running.

Table 2 provides an overview of the running economy characteristics associated with the outcomes of concurrent HIIT and MICT interventions. Running economy was evaluated by measuring the oxygen cost associated with each condition at a specific running speed. Statistically significant improvements in RE were observed in at least one of the nine studies. Five studies assessed RE at a single speed, while four others evaluated RE at multiple speeds. Additionally, maximal oxygen uptake and blood lactate concentration were included as secondary measures to assess other factors influencing RE.

Assessment of Quality

Figure 2 shows a statistical chart illustrating the proportion of projects evaluated using different methods. Among the nine articles considered, the blind method posed the highest risk, as the experimental intervention necessitated participants to sign informed consent, leading to significant bias (Figure 2A). Three studies exhibited low risk of bias and high quality, while the remaining six studies had moderate risk of bias.

Primary Outcomes

Running Economy

This systematic review included nine randomized controlled trials (RCTs) that assessed running economy (RE) at velocities ranging from 7.00 to 16.00 km/h. Due to variability in gender, age, units of measurement, and test intensity, a random effects model was applied. The lactate threshold refers to the point at which blood lactate

levels rise nonlinearly. The first lactate threshold (LT1) occurs at approximately 2 $\text{mmol}\cdot\text{L}^{-1}$, while the second lactate threshold (LT2) is observed at approximately 4 $\text{mmol}\cdot\text{L}^{-1}$ (Seiler and Kjerland, 2006). The ventilation threshold exhibits a similar pattern, with the first ventilation threshold (VT1) corresponding to the LT1 (approximately 75% $\text{VO}_{2\text{max}}$) and the second ventilation threshold (VT2) aligning with the LT2 (approximately 85% $\text{VO}_{2\text{max}}$) (Keir et al., 2022). Based on the research conducted by Casado et al. (2022), this study categorized exercise intensity into three commonly utilized training zones for subgroup analysis: Zone 1 (Z1) corresponded to speeds below the first ventilatory or lactate threshold (approximately 70% $\text{VO}_{2\text{max}}$), Zone 2 (Z2) included speeds between the first and second thresholds (approximately 75%–85% $\text{VO}_{2\text{max}}$), and Zone 3 (Z3) encompassed speeds exceeding the second threshold (above 85% $\text{VO}_{2\text{max}}$) (Casado et al., 2022; Stögg and Sperlich, 2015). Most of the included subjects in this study had oxygen uptake levels between 50 and 60, and only in two studies participants had oxygen uptake levels around 35 and 70, respectively, which may have impacted the results. Therefore, we conducted sensitivity analysis on these two studies and found that their results did not significantly affect the overall results of the study. In Z1, the heterogeneity among studies was low ($I^2 = 7\%$, $p = 0.38$). The standardized mean difference (SMD) was 0.40, with a 95% confidence interval of [0.04, 0.76], indicating statistical significance ($Z = 2.16$, $p < 0.05$). These results suggest that HIIT was more effective than MICT at improving RE at speeds within Z1. In Z2, there was minimal heterogeneity among the studies ($I^2 = 34\%$, $p = 0.19$). The standardized mean difference (SMD) was 0.68, with a 95% confidence interval of [0.13, 1.24], showing statistical significance ($Z = 2.40$, $p < 0.05$). The results indicated that subjects following HIIT exhibited better running economy and consumed less oxygen compared to those following MICT at speeds within Z2. In Z3, where speeds exceeded the lactate threshold, there was no intra-group heterogeneity ($I^2 = 40\%$, $p > 0.1$). The standardized mean difference (SMD) was 0.14, with a 95% confidence interval of [-0.60, 0.88]. There was no statistical significance in training results within Z3 ($Z = 0.37$, $p > 0.05$), which indicates that there was no significant difference in subjects' running economy between HIIT and MICT at speeds

exceeding the lactate threshold.

The heterogeneity between groups was low ($I^2 = 24\%$, $p > 0.1$). The standardized mean difference (SMD) between groups was 0.44, with a 95% confidence interval of [0.15, 0.72], indicating statistical significance ($Z = 3.01$, $p < 0.05$). This indicates that both HIIT and MICT had a greater effect on RE, with the control group showing a larger total effect size than the experimental group. This also suggests that the control group consumed more oxygen under similar conditions, while the experimental group consumed less oxygen. Thus, it can be stated that HIIT was more effective in improving RE.

Maximal Oxygen Uptake

Changes in maximal oxygen uptake ($VO_{2\max}$) were compared between HIIT and MICT, before and after the experiment. The heterogeneity among the groups was low ($I^2 = 38\%$, $p = 0.12$), which justified the use of a fixed-effects model. In general, when the I^2 value is below 50% (or the commonly used threshold is 25%–50%), the heterogeneity is small, the results tend to be consistent, and the use of fixed-effect models is a suitable choice. The mean difference (MD) was 2.48, with a 95% confidence interval of [1.61, 3.34], indicating statistical significance ($Z = 5.60$, $p < 0.05$). The findings revealed significant differences in $VO_{2\max}$ between the two training regimens, with MICT showing a greater improvement compared to HIIT.

Blood Lactate Variables

The blood lactate levels were measured at the velocity associated with $VO_{2\max}$ in three studies, while fixed blood lactate levels were assessed in one study. The effects of HIIT and MICT on blood lactate levels were evaluated before and after the intervention. There was no heterogeneity among the groups ($I^2 = 0\%$, $p > 0.1$), and a fixed-effect model was employed for the analysis. The research summary indicated a mean difference (MD) of 0.15, with a 95% confidence interval of [-0.28, -0.02], which was statistically significant ($Z = 2.20$, $p < 0.05$). The results indicated that lactic acid production in HIIT was lower than that in MICT at the same intensity following training. Additionally, lactic acid tolerance was improved.

Analysis of Publication Bias

A funnel plot was created to assess running

economy as the primary outcome measure. The funnel plot was visually inspected, and it appeared symmetric, suggesting that the included studies were fairly distributed around the pooled effect size. This symmetry typically indicates the absence of significant publication bias, as smaller studies are equally likely to show both positive and negative results. The Egger's test for asymmetry yielded no significant results ($p > 0.05$), and further statistical tests indicated that the distribution of study effects was likely unbiased. However, it is important to note that while no significant publication bias was detected, small studies with null or negative results may still be underreported, potentially leading to some unobserved bias. Overall, the evidence suggests that publication bias did not significantly affect the findings of this meta-analysis, although caution should always be taken when interpreting results from small studies with non-significant outcomes.

Discussion

The objective of this systematic review was to critically evaluate the existing literature to determine the effects of HIIT and MICT on running economy (RE). The initial findings revealed that HIIT significantly reduced the oxygen cost of running compared to MICT in endurance runners. Furthermore, the effects of different training interventions on maximal oxygen uptake indicated that MICT was more effective in improving $VO_{2\max}$. Finally, training had a significant impact on blood lactate concentration. HIIT was particularly effective as it enabled the body to adapt to lactate more efficiently than most of other forms of exercise. Individuals who engaged in HIIT produced less lactate at similar exercise intensities and demonstrated higher tolerance to lactate.

Table 1. Characteristics of training intervention of considered studies.

Study G	Subjects' characteristics				VO _{2max} (ml·kg ⁻¹ ·min ⁻¹)			Training intervention						
	n(M/F)	Age	H	W	BMI	Pre	post	PL	D	Fq	TW	exercise	Load	rest
Reuter et al. (2024)	HIIT 15 (7M; 8F)	45.9 ± 9	171.2 ± 5.3	75 ± 12	25.4 ± 3.5	34.0 ± 5.2	37.3 ± 4.9	MT	26	3	84 min	4 × 4 min	95% HR _{max}	3 min
	MICT 16 (6M; 10F)	46.9 ± 7.8	170.6 ± 9.9	75 ± 16	25.5 ± 3.3	34.9 ± 3.6	35.3 ± 4	MT	26	3	150 min	50 min × 1 session	55%HR _{reserve}	
Rago et al. (2022)	HIIT 11 (M)	22.0 ± 1.8	184.9 ± 5.4	76.9 ± 5.4	22.49 ± 1.58	60.82 ± 2.66	62.56 ± 3.42	WT	10	2	360 s	6–12 × 30-s sprints	~95%VO ₂	3 min
	MICT 11 (M)	21.9 ± 2.3	182.4 ± 7.9	73.0 ± 6.1	21.94 ± 1.83	60.56 ± 3.10	61.1 ± 3.81	WT	102		20–40 min	2 × 1 week of low-intensity running	70% VO _{2max}	
Festa et al. (2020)	HIIT 19 (15M; 4F)	43.2 ± 8.4	175.2 ± 5.9	72.0 ± 7.7	23.46 ± 0.02	52.9 ± 8.1	53.6 ± 4.8	MT	8	4	3.74 ± 0.38 h	2 high Intensities running		
	MICT 19 (16M; 3F)	39.4 ± 8.5	172.5 ± 4.3	70.9 ± 10.1	23.83 ± 0.14	53.4 ± 8.3	53.2 ± 1.9	MT	8	4	3.1 ± 0.25 h	moderate intensity		
Jarstad and Mamen (2019)	HIIT 7 (4M; 3F)	28 ± 5	NR	78.4 ± 11	23.9 ± 2.3	50.6 ± 5.4	53.5 ± 6.9	MT	10	3	60 min	20-min strenuous exhausting run	83% VO _{2max}	NR
	MICT 7 (4M; 4F)	28 ± 5	NR	76.3 ± 10.4	24.0 ± 2.7	50.9 ± 7.1	53.3 ± 7.2	MT	10	3	120 min	40-min run at ~80% HR _{max}	72% VO _{2max}	NR
Silva et al. (2017)	HIIT 8 (M)	35 ± 6	172.5 ± 4.1	70.5 ± 4.6	NR	54.5 ± 8.1	57.1 ± 6.4	MT	4	2	265 ± 67 s	high-intensity interval training twice weekly		NR
	MICT 8 (M)	32 ± 9	172.8 ± 9	70.2 ± 11.3		56.6 ± 7.3	56.9 ± 7.6	MT	4	2		regular endurance training		
González-Mohino et al. (2016)	HIIT 6		NR			57.71 ± 8.92	56.2 ± 6.685	MT	6	3	29 min	Interval training 95–110% of MAS	95–110% of MAS	1min
	MICT 5					55.48 ± 8.45	51.66 ± 6.68	MT	6	3	60 min	Interval training 70–75% of MAS	70–75% of MAS	
Gojanovic et al. (2015)	HIIT 6 (M)	31.7 ± 5.2	182 ± 10	77.3 ± 5.2	23.4 ± 2	54.1 ± 8.3	55.1 ± 7.6	MT	4	2	534 s	4–5 intervals run at 100% of velocity to exhaustion	vVO _{2max}	267 ± 68 s
	MICT 5 (M)	38.4 ± 9.7	178 ± 4	75.9 ± 3.6	24.1 ± 1.6	57.6 ± 2.5	59.6 ± 2	MT	4	2	368 s			
Enoksen et al. (2011)	HIIT 9 (M)		NR			70.2 ± 2.7	71.4 ± 2.4	HT	10	6	NR	intensive workouts per week at 82–92% of HR _{max}	82% HR _{max}	NR
	MICT 10 (M)					70.4 ± 3.8	69.2 ± 3.6	HT	10	6	NR	performed at 65–82% of HR _{max}	65% HR _{max}	
Litleskare et al. (2020)	HIIT 13 (5M; 8F)	25 ± 1	175 ± 3	71.2 ± 4.1	24.0 ± 0.8	50.5 ± 1.6	53.3 ± 1.5	MT	8	3	150 s	5–10 per session 30-s sprints at near maximal effort	85% HR _{peak}	3 min
	MICT 12 (4M; 8F)	25 ± 1	173 ± 2	72.6 ± 3.8	23.6 ± 0.9	47.9 ± 1.5	49.7 ± 1.5	MT	8	3	150 min	70%HR _{peak} training sessions 30 min up to 60 min	70% HR _{peak}	

HIIT: high-intensity interval training, MICT: moderate-intensity continuous training, G: group, n: sample size, F: female, M: male, Age (years), H: body height (cm), W: body weight (kg), BMI: body mass index, pre untrained, post trained, VO₂: maximal oxygen uptake, PL: performance level, D: duration (weeks), Fq: frequency (session/week), TW: Weekly training hours, HT: highly trained, MT: moderately trained, WT: well trained, NR: not reported, MAS: Velocity at VO₂, V: Velocity, HR_{max}: maximum heart rate, HR_{peak}: peak heart rate, HR_{reserve}: maximum heart rate reserve, s: seconds, min: minute

Table 2. Running economy of studies included in the meta-analysis.

Study	G	N	running economy						
			measurement	speed reported	Relative to AT	Relative to VO _{2max} (%)	mean pre (SD)	mean post (SD)	outcomes
Reuter et al. (2024)	HIIT	15(7M; 8F)	ml·kg ⁻¹ ·min ⁻¹	7	<AT	70% VO _{2max}	157 ± 75	152 ± 73	①②③
	MICT	16(6M; 10F)	ml·kg ⁻¹ ·min ⁻¹	7	<AT	70% VO _{2max}	161 ± 50	150 ± 47	①②③
	HIIT	11(M)	ml·kg ⁻¹ ·min ⁻¹	10	<AT		224.00 ± 12.84	218.16 ± 11.82	①②
Rago et al. (2022)	MICT	11(M)	ml·kg ⁻¹ ·min ⁻¹	12	≤AT		219.55 ± 13.97	214.74 ± 11.34	
			ml·kg ⁻¹ ·min ⁻¹	14	≤AT		215.17 ± 12.29	216.47 ± 9.70	
			ml·kg ⁻¹ ·min ⁻¹	10	<AT		211.31 ± 18.79	215.29 ± 18.24	①②
			ml·kg ⁻¹ ·min ⁻¹	12	≤AT		202.84 ± 18.83	206.46 ± 18.75	
			ml·kg ⁻¹ ·min ⁻¹	14	≤AT		207.47 ± 20.58	210.94 ± 20.71	
Festa et al. (2020)	HIIT	19(15M; 4F)	ml·kg ⁻¹ ·min ⁻¹	8.9 ± 0.2	<AT		226.3 ± 35.2	214.3 ± 33.0	①②
	MICT	19(16M; 3F)	ml·kg ⁻¹ ·min ⁻¹	8.9 ± 0.2	<AT		231.8 ± 9.1	211.6 ± 6.3	①②
Jarstad and Mamen (2019)	HIIT	7(4M; 3F)	ml·kg ^{-0.75} ·m ⁻¹	8.9 ± 1.2	<AT	70% VO _{2max}	0.721 ± 0.064	0.694 ± 0.064	①②③
	MICT	7(4M; 4F)	ml·kg ^{-0.75} ·m ⁻¹	9.6 ± 0.8	<AT	70% VO _{2max}	0.664 ± 0.054	0.685 ± 0.058	①②③
Silva et al. (2017)	HIIT	8(M)	ml·kg ⁻¹ ·min ⁻¹	12	≤AT		43.1 ± 3.5	40.7 ± 4.3	①②
	MICT	8(M)	ml·kg ⁻¹ ·min ⁻¹	12	≤AT		40.9 ± 4.7	41.2 ± 4.4	①②
	HIIT	6	ml·kg ⁻¹ ·min ⁻¹	10.0 ± 0.88	<AT	60% VO _{2max}	37.55 ± 4.92	38.55 ± 4.23	①②
González-Mohino et al. (2016)	MICT	5	ml·kg ⁻¹ ·min ⁻¹	14.53 ± 1.18	≤AT	80% VO _{2max}	48.3 ± 5.41	46.81 ± 6.44	
			ml·kg ⁻¹ ·min ⁻¹	16.34 ± 1.32	>AT	90% VO _{2max}	52.6 ± 6.96	51.8 ± 5.19	
			ml·kg ⁻¹ ·min ⁻¹	9.84 ± 1.09	<AT	60% VO _{2max}	38.46 ± 2.46	32.66 ± 4.54	①②
			ml·kg ⁻¹ ·min ⁻¹	13.12 ± 1.45	≤AT	80% VO _{2max}	45.62 ± 3.77	43.22 ± 3.88	
			ml·kg ⁻¹ ·min ⁻¹	14.76 ± 1.63	>AT	90% VO _{2max}	50.86 ± 5.68	46.86 ± 5.77	
Gojanovic et al. (2015)	HIIT	6(M)	ml·kg ⁻¹ ·min ⁻¹	14	≤AT		47.1 ± 4	45.9 ± 4.6	①②
	MICT	5(M)	ml·kg ⁻¹ ·min ⁻¹	14	≤AT		48.4 ± 4.9	47.3 ± 3.7	①②
Enoksen et al. (2011)	HIIT	9	ml·kg ⁻¹ ·min ⁻¹	13	≤AT		51.1 ± 3.8	48.7 ± 3.0	①②③
			ml·kg ⁻¹ ·min ⁻¹	14.5	≤AT		56.9 ± 3.3	55.1 ± 3.2	
			ml·kg ⁻¹ ·min ⁻¹	16	>AT		62.0 ± 3.0	60.4 ± 2.5	
			ml·kg ⁻¹ ·min ⁻¹	13	≤AT		49.6 ± 2.3	47.5 ± 1.7	①②③
			ml·kg ⁻¹ ·min ⁻¹	14.5	≤AT		55.3 ± 2.7	52.8 ± 2.7	
			ml·kg ⁻¹ ·min ⁻¹	16	>AT		61.4 ± 2.2	59.0 ± 2.6	
			ml·kg ⁻¹ ·min ⁻¹	13(5M; 8F)	<AT	50% VO _{2max}	201 ± 10	199 ± 10	①②③
			ml·kg ⁻¹ ·min ⁻¹	10.4 ± 0.3	≤AT	80% VO _{2max}	237 ± 4	231 ± 4	
Litleskare et al. (2020)	MICT	12(4M; 8F)	ml·kg ⁻¹ ·min ⁻¹	6.2 ± 0.2	<AT	50% VO _{2max}	213 ± 16	186 ± 10	①②③
			ml·kg ⁻¹ ·min ⁻¹	10.1 ± 0.3	≤AT	80% VO _{2max}	232 ± 6	223 ± 6	

HIIT: high-intensity interval training, MICT: moderate-intensity continuous training, G: group, N: sample size, F: female, M: male, AT: anaerobic threshold, ①Running Economy, ②Maximal Oxygen Uptake, ③Blood Lactate




Figure 1. Flow diagram of the search strategy.




Figure 2. Assessment of quality. Note: In Figure 2(B), the "+" symbol represents low risk of bias, whereas the "-" symbol indicates high risk of bias





Figure 3. Forest plot for running economy.

Figure 4. Forest plot for VO_{2max}.




Figure 5. Forest plot for blood lactate concentration.




Figure 6. The risk of bias funnel plot.

The objective of this systematic review was to critically evaluate the existing literature to determine the effects of HIIT and MICT on running economy (RE). The initial findings revealed that HIIT significantly reduced the oxygen cost of running compared to MICT in endurance runners. Furthermore, the effects of different training interventions on maximal oxygen uptake indicated that MICT was more effective in improving $VO_{2\max}$. Finally, training had a significant impact on blood lactate concentration. HIIT was particularly effective as it enabled the body to adapt to lactate more efficiently than most of other forms of exercise. Individuals who engaged in HIIT produced less lactate at similar exercise intensities and demonstrated higher tolerance to lactate.

Our initial findings showed that, among endurance runners, HIIT led to significantly reduced oxygen cost of running compared to MICT (Wang et al., 2012). Previous studies have indicated that different training interventions can have distinct effects on running economy (Barnes and Kilding, 2015; Faelli et al., 2021; Franch et al., 1998). Compared to MICT interventions, HIIT intervals of 1 min or less did not demonstrate a significant difference in oxygen costs. However, intervals longer than one minute led to a significant improvement in maximal oxygen uptake (Hov et al., 2023). Increasing the work interval from 30 to 60 s led to an increase in maximal oxygen uptake, despite shorter total training duration (Wen et al., 2019). Doubling the interval duration (2 min versus 1 min) resulted in a substantial increase in anaerobic glycolytic energy release, along with an even greater enhancement in oxygen availability (Vuorimaa et al., 2000). HIIT primarily acts on fast-twitch muscle fibers through the arrangement of interval training duration and recovery time (interval training with duration of ≥ 2 min being particularly effective for aerobic stimulation); MICT primarily targets slow muscle fibers, through continuous training, it increases the number of mitochondria and the expression of related proteases in slow muscles (Skelly et al., 2021). It not only enhances the functions of the autonomic nervous system and parasympathetic nervous systems (Clemente-Suárez and Arroyo-Toledo, 2018), but also induces lactic acid accumulation, activates antioxidant enzyme systems, and improves glycolytic capacity through rapid muscle contractions and exertion (Peake et

al., 2014). Additionally, HIIT enhances muscle and mitochondrial size (Estes et al., 2017; Ruegsegger et al., 2023), increases hemoglobin levels, and improves blood flow-mediated vasodilation (Mandić et al., 2022; Sawyer et al., 2016). These adaptations contribute to the overall efficiency of both aerobic and anaerobic energy systems (Bogdanis et al., 2013; Buchheit and Laursen, 2013). MICT can increase muscle capillary density, improve myocardial perfusion, enhance blood flow-mediated dilation, and facilitate oxygen delivery (Cocks et al., 2016; Daussin et al., 2008). These effects are beneficial for energy supply during exercise and contribute to the body's long-term resistance to fatigue (Wang et al., 2021). The spring-mass model is an important factor associated with RE, where the body's bounce on the ground is counteracted by the spring-like behavior of the supporting leg. Mechanical energy is stored in the muscles, tendons, and ligaments acting across the joints (Saunders et al., 2004). HIIT can have greater impact on muscle elasticity than MICT, the faster the speed, and the higher the stiffness in the lower limb muscles. The greater the stiffness, the more beneficial it is for runners to consume less oxygen (Liu et al., 2022). Compared with MICT, HIIT brings greater physical improvements and adaptive changes over a short period of time, thereby enhancing RE of runners (Skovgaard et al., 2018a). According to our meta-analysis, HIIT has more beneficial effects on RE at intensities near Z2 compared to those near Z3. When the velocity exceeds Z2, the effect size becomes non-significant (Figure 3). Training at an intensity close to Z2 may elicit a combination of molecular signals generated by both high-volume and high-intensity training. This combination of signals provides greater stimulation and has a more significant impact on the body's adaptation, which is more conducive to improving RE (Skovgaard et al., 2018b). Training in Z3 primarily stimulates the body anaerobically, and although this intensity can also stimulate the body's aerobic metabolism to a certain extent, the effect is inferior to the former. This finding is consistent with research indicating that when lactate concentration is greater than 4 mmol/L, the enhancement in running economy diminishes (Zinner et al., 2018). Furthermore, the impact on running economy continues to decline as exercise intensity increases (Munoz et al., 2015). Among the different runners,

we found that the improvement in running economy within the Z3 was reduced for well-trained and recreational runners, with the effect being particularly diminished for recreational runners. Repeated training in Z3 may result in muscle damage, fatigue, and decreased parasympathetic nerve activity (Kingsley et al., 2005; Nuutila et al., 2022). For runners, training intensity should be kept close to Z2 to optimize running economy, especially for low- and medium-level athletes. Overall, our findings indicate that HIIT at or below the lactate threshold during the Z2 phase is essential for enhancing running economy.

It is widely acknowledged that maximal oxygen uptake is one of the most critical factors for successful distance running (Parmar et al., 2021). Current studies have shown that both HIIT and MICT interventions can enhance maximal oxygen uptake (Mølmen et al., 2025). Our meta-analysis found that MICT had a greater effect on peak oxygen uptake than HIIT, which is consistent with a recent review of related studies (Guo et al., 2022; Liang et al., 2024). However, other scholars have expressed an opposing viewpoint. Some researchers pointed out that HIIT was particularly effective in improving $VO_{2\max}$ during the initial stages of training for individuals with lower fitness levels (Mølmen et al., 2025). However, improvements in ventilatory thresholds above the second ventilatory improvements were only observed in well-trained endurance runners (Seiler et al., 2007; Tanji and Nabekura, 2019). HIIT can cause muscle damage and pain, increase cellular damage, lipid peroxidation, inflammatory response, and adversely affect muscle cell structure and the function of contractile proteins (Kingsley et al., 2005; Leite et al., 2023). HIIT for recreational runners may increase their risk of muscle injuries (Leite et al., 2023; Nuutila et al., 2022). Additionally, it can lead to decreased adherence to training, particularly in unsupervised settings (Ekkekakis and Biddle, 2023; Santos et al., 2023). In contrast, MICT is relatively effective in mitigating these risks and enhancing runners' compliance with their training regimens (Ekkekakis and Biddle, 2023; Santos et al., 2023). Our research indicates that MICT is a more effective way to develop aerobic endurance than HIIT. For most runners, particularly beginners and those who are untrained, utilizing MICT to

enhance aerobic capacity still has its significant advantages.

Blood lactate concentration is an important indicator of an individual's exercise intensity. Distance running performance can be partially predicted by blood lactate concentration at a specific running speed (Farrell et al., 1979; Fay et al., 1989; Yoshida et al., 1993). This research included four studies on blood lactate concentration, which demonstrated that HIIT was more effective in adapting blood lactate concentration than MICT. This may be attributed to an increased muscle oxidative capacity and associated changes in motor unit recruitment patterns, as well as muscle adaptations, enhanced glycolytic capacity, increased oxidative activity, and alterations in lactate transport (Denadai et al., 2006; Joyner and Coyle, 2008). These benefits include enhanced recovery of creatine phosphate and increased mitochondrial biogenesis, improved skeletal muscle buffering capacity, enhanced ventilation, and elevated lactate threshold (Driller et al., 2009; Esfarjani and Laursen, 2007; Hoogeveen, 2000; Weston et al., 1997). Additionally, there is an increased capacity to engage larger volumes of muscle mass (Creer et al., 2004) and a greater ability to oxidize fat relative to carbohydrates (Atakan et al., 2022). Compared with MICT, HIIT can delay lactic acid accumulation in the blood and postpone muscle fatigue during high-intensity exercise, resulting in improved physical performance and improved training outcomes.

Both funnel plot and the Egger's test showed no obvious publication bias for the included studies. These studies exhibited diversity in terms of the study sites, populations, and designs, which reduced the likelihood of publication bias to some extent. Since different study settings and designs produce different effect size estimates, it is difficult to guarantee a consistent positive outcome preference across such diverse studies if publication bias exists. We found some differences in the maximum oxygen uptake of Reuter et al. (2024) and Enoksen et al.'s (2011) studies, which were about 35 and 70, respectively. This difference may bias the results of the studies, thus sensitivity analysis was conducted one by one. It was found that the inclusion and exclusion of these two articles had no significant effect on the results of the study. We weighted each study based

on its quality, sample size, and other factors. Despite potential bias in some studies (e.g., differences in oxygen uptake between Reuter et al. (2024) and Enoksen et al.'s (2011) studies), sensitivity analysis showed that including or excluding these studies did not significantly affect the final results. Despite some bias, the weighting method and analysis results remained stable and reliable.

The limitations of this systematic review include methodological and statistical heterogeneity among the included studies, with some exhibiting risk of bias due to methodological weaknesses, such as lack of blinding in intervention studies. Varying measurement methods may have contributed to heterogeneous results, and although subgroup analysis addressed some sources of heterogeneity, individual-level moderators remained. Furthermore, the studies did not account for other influencing factors, such as environmental conditions, sleep, diet, and psychological stress, which can affect running economy. For instance, acute sleep deprivation can impair performance (Craven et al., 2022), leading to an increased heart rate and minute ventilation during exercise (Mougin et al., 1991). Poor sleep can also deteriorate athletic performance, increasing energy cost during exercise (Vitale et al., 2019). Additionally, moderate carbohydrate intake before and after exercise can improve endurance and glycogen reserves (Flockhart and Larsen, 2024; Khanna and Manna, 2005), while proper protein supplementation aids muscle synthesis and repair (Phillips and Van Loon, 2011). Dietary intake of carbohydrates and fats influences fuel utilization and running economy (Maunder et al., 2021), with low carbohydrate (<20%) and high fat (>60%) intake promoting fat oxidation and improved submaximal running economy (Che et al., 2021). Psychological stress can also impact performance by altering hormone levels, such as cortisol and adrenaline (Rimmele et al., 2009; Schommer et al., 2003). Pre-match anxiety can elevate the heart rate and blood pressure, impairing energy utilization and performance (Nibbeling et al., 2012; Webb et

al., 2008). Lastly, the review was restricted to English-language studies and to the sport of running, which may limit the generalizability of the findings. Therefore, caution is needed when interpreting the results, and future research should explore running economy in other sports and consider papers published in other languages for broader applicability.

HIIT has gained popularity, but studies comparing its long-term effects with MICT show no significant superiority. While short-term compliance with HIIT is higher (Jung et al., 2015), long-term adherence drops significantly, with dropout rates higher than MICT (Roy et al., 2018). One study found adherence dropping from 82% to 29% at 12 months. During a 12-month follow-up, MICT participants were more likely to complete training (Taylor et al., 2020). HIIT also carries higher risks, with the incidence of acute cardiac events nearly six times higher (Rognmo et al., 2012), and muscle function impairment linked to elevated blood lactate concentration (Thomas et al., 2004). The injury risk associated with different interval duration and intensities during HIIT may lead to musculoskeletal pain, especially at higher running speeds (Leite et al., 2023; Wiewelhove et al., 2016). Therefore, the high intensity, high risk of injury, and low long-term adherence of HIIT make it more challenging for long-term sustainability in mass fitness participants.

Conclusions

The results of this study suggest that HIIT is a more effective method for improving running economy and delaying lactate accumulation, as well as enhancing lactate tolerance, compared to MICT in long-distance runners. By training close to the lactate threshold (Zone 2), HIIT facilitates greater improvements in running economy and postpones lactate accumulation in the bloodstream. However, improvements in maximal oxygen uptake ($VO_{2\max}$) were significantly more pronounced with MICT than with HIIT. Further high-quality studies are needed to investigate these effects in greater depth in the future.

Author Contributions: Conceptualization: Y.F.; methodology: D.L.; software: D.L. and Y.L.; validation: Y.F., D.L. and Y.L.; formal analysis: Y.L.; investigation: D.L.; resources: Y.F.; data curation: D.L.; writing—original draft preparation: Y.L.; writing—review & editing: Y.F.; visualization: D.L.; supervision: D.T.; project administration: Y.F.; funding acquisition: D.T. All authors have read and agreed to the published version of the manuscript.

ORCID iD:

Donghui Tang: <https://orcid.org/0009-0002-9097-007X>

Funding Information: This research received no external funding.

Institutional Review Board Statement: Not applicable.

Informed Consent: Not applicable.

Conflicts of Interest: The authors declare no conflict of interest.

Received: 09 September 2024

Accepted: 21 May 2025

References

Atakan, M. M., Guzel, Y., Shrestha, N., Kosar, S. N., Grgic, J., Astorino, T. A., Turnagol, H. H., & Pedišić, Z. (2022). Effects of high-intensity interval training (HIIT) and sprint interval training (SIT) on fat oxidation during exercise: a systematic review and meta-analysis. *British Journal of Sports Medicine*, 56(17), 988–996. <https://doi.org/10.1136/bjsports-2021-105181>

Barnes, K. R., & Kilding, A. E. (2015). Strategies to improve running economy. *Sports Medicine*, 45(1), 37–56. <https://doi.org/10.1007/s40279-014-0246-y>

Baroni, B. M., Rodrigues, R., Franke, R. A., Geremia, J. M., Rassier, D. E., & Vaz, M. A. (2013). Time course of neuromuscular adaptations to knee extensor eccentric training. *International Journal of Sports Medicine*, 34(10), 904–911. <https://doi.org/10.1055/s-0032-1333263>

Black, M. I., Handsaker, J. C., Allen, S. J., Forrester, S. E., & Folland, J. P. (2018). Is there an optimal speed for economical running? *International Journal of Sports Physiology and Performance*, 13(1), 75–81. <https://doi.org/10.1123/ijsspp.2017-0015>

Bogdanis, G. C., Stavrinou, P., Fatouros, I. G., Philippou, A., Chatzinikolaou, A., Draganidis, D., Ermidis, G., & Maridaki, M. (2013). Short-term high-intensity interval exercise training attenuates oxidative stress responses and improves antioxidant status in healthy humans. *Food and Chemical Toxicology: An International Journal Published for the British Industrial Biological Research Association*, 61, 171–177. <https://doi.org/10.1016/j.fct.2013.05.046>

Buchheit, M., & Laursen, P. B. (2013). High-intensity interval training, solutions to the programming puzzle. Part II: anaerobic energy, neuromuscular load and practical applications. *Sports Medicine*, 43(10), 927–954. <https://doi.org/10.1007/s40279-013-0066-5>

Casado, A., González-Mohín, F., González-Ravé, J. M., & Foster, C. (2022). Training periodization, methods, intensity distribution, and volume in highly trained and elite distance runners: a systematic review. *International Journal of Sports Physiology and Performance*, 17(6), 820–833. <https://doi.org/10.1123/ijsspp.2021-0435>

Che, K., Qiu, J., Yi, L., Zou, M., Li, Z., Carr, A., Snipe, R. M. J., & Benardot, D. (2021). Effects of a short-term “fat adaptation with carbohydrate restoration” diet on metabolic responses and exercise performance in well-trained runners. *Nutrients*, 13(3), 1033. <https://doi.org/10.3390/nu13031033>

Clemente-Suárez, V. J., & Arroyo-Toledo, J. J. (2018). The use of autonomic modulation device to control training performance after High-Intensity interval training program. *Journal of Medical Systems*, 42(3), 47. <https://doi.org/10.1007/s10916-018-0907-8>

Cocks, M., Shaw, C. S., Shepherd, S. O., Fisher, J. P., Ranasinghe, A., Barker, T. A., & Wagenmakers, A. J. (2016). Sprint interval and moderate-intensity continuous training have equal benefits on aerobic capacity, insulin sensitivity, muscle capillarisation and endothelial eNOS/NAD(P)H oxidase protein ratio in obese men. *Journal of Physiology*, 594(8), 2307–2321. <https://doi.org/10.1113/jphysiol.2014.285254>

Craven, J., McCartney, D., Desbrow, B., Sabapathy, S., Bellinger, P., Roberts, L., & Irwin, C. (2022). Effects of acute sleep loss on physical performance: a systematic and meta-analytical review. *Sports Medicine*, 52(11), 2669–2690. <https://doi.org/10.1007/s40279-022-01706-y>

Creer, A. R., Ricard, M. D., Conlee, R. K., Hoyt, G. L., & Parcell, A. C. (2004). Neural, metabolic, and performance adaptations to four weeks of high intensity sprint-interval training in trained cyclists. *International Journal of Sports Medicine*, 25(2), 92–98. <https://doi.org/10.1055/s-2004-819945>

Daussin, F. N., Zoll, J., Dufour, S. P., Ponsot, E., Lonsdorfer-Wolf, E., Doutreleau, S., Mettauer, B., Piquard, F., Geny, B., & Richard, R. (2008). Effect of interval versus continuous training on cardiorespiratory and mitochondrial functions: relationship to aerobic performance improvements in sedentary subjects. *American Journal of Physiology Regulatory Integrative and Comparative Physiology*, 295(1), R264–R272. <https://doi.org/10.1152/ajpregu.00875.2007>

Del Vecchio, A., Casolo, A., Negro, F., Scorceti, M., Bazzucchi, I., Enoka, R., Felici, F., & Farina, D. (2019). The increase in muscle force after 4 weeks of strength training is mediated by adaptations in motor unit recruitment and rate coding. *Journal of Physiology*, 597(7), 1873–1887. <https://doi.org/10.1113/JP277250>

Denadai, BS., Ortiz, MJ., Greco, CC., & Mello, MT. (2006). Interval training at 95% and 100% of the velocity at $VO_{2\text{max}}$: effects on aerobic physiological indexes and running performance. *Applied Physiology, Nutrition, and Metabolism*, 31(6), 737–743. <https://doi.org/10.1139/h06-080>

Dohlmann, T. L., Hindsø, M., Dela, F., Helge, J. W., & Larsen, S. (2018). High-intensity interval training changes mitochondrial respiratory capacity differently in adipose tissue and skeletal muscle. *Physiological Reports*, 6(18), e13857. <https://doi.org/10.14814/phy2.13857>

Dolan, C., Quiles, J. M., Goldsmith, J. A., Mendez, K. M., Klemp, A., Robinson, Z. P., Pelland, J. C., Coccia, C., & Zourdos, M. C. (2024). The effect of time-equated concurrent training programs in resistance-trained men. *Journal of Human Kinetics*, 91, 87–103. <https://doi.org/10.5114/jhk/185637>

Driller, M. W., Fell, J. W., Gregory, J. R., Shing, C. M., & Williams, A. D. (2009). The effects of high-intensity interval training in well-trained rowers. *International Journal of Sports Physiology and Performance*, 4(1), 110–121. <https://doi.org/10.1123/ijspp.4.1.110>

D'Souza, R. F., Woodhead, J. S. T., Zeng, N., Blenkiron, C., Merry, T. L., Cameron-Smith, D., & Mitchell, C. J. (2018). Circulatory exosomal miRNA following intense exercise is unrelated to muscle and plasma miRNA abundances. *American Journal of Physiology. Endocrinology and Metabolism*, 315(4), E723–E733. <https://doi.org/10.1152/ajpendo.00138.2018>

Ekkekakis, P., & Biddle, S. J. H. (2023). Extraordinary claims in the literature on high-intensity interval training (HIIT): IV. Is HIIT associated with higher long-term exercise adherence? *Psychology of Sport and Exercise*, 64, 102295. <https://doi.org/10.1016/j.psychsport.2022.102295>

Enoksen, E., Shalfawi, S. A., & Tønnessen, E. (2011). The effect of high- vs. low-intensity training on aerobic capacity in well-trained male middle-distance runners. *Journal of Strength and Conditioning Research*, 25(3), 812–818. <https://doi.org/10.1519/JSC.0b013e3181cc2291>

Esfarjani, F., & Laursen, P. B. (2007). Manipulating high-intensity interval training: effects on $VO_{2\text{max}}$, the lactate threshold and 3000 m running performance in moderately trained males. *Journal of Science and Medicine in Sport*, 10(1), 27–35. <https://doi.org/10.1016/j.jsams.2006.05.014>

Estes, R. R., Malinowski, A., Piacentini, M., Thrush, D., Salley, E., Losey, C., & Hayes, E. (2017). The effect of high intensity interval run training on cross-sectional area of the vastus lateralis in untrained college students. *International Journal of Exercise Science*, 10(1), 137–145. <https://doi.org/10.70252/ULIC9262>

Faelli, E., Panascì, M., Ferrando, V., Bisio, A., Filipas, L., Ruggeri, P., & Bove, M. (2021). The effect of static and dynamic stretching during warm-up on running economy and perception of effort in recreational endurance runners. *International Journal of Environmental Research and Public Health*, 18(16), 8386. <https://doi.org/10.3390/ijerph18168386>

Farrell, P. A., Wilmore, J. H., Coyle, E. F., Billing, J. E., & Costill, D. L. (1979). Plasma lactate accumulation and distance running performance. *Medicine and Science in Sports*, 11(4), 338–344.

Fay, L., Londeree, B. R., LaFontaine, T. P., & Volek, M. R. (1989). Physiological parameters related to distance running performance in female athletes. *Medicine and Science in Sports and Exercise*, 21(3), 319–324. <https://doi.org/10.1249/000005768-198906000-00016>

Festa, L., Tarperi, C., Skroce, K., La Torre, A., & Schena, F. (2020). Effects of different training intensity distribution in recreational runners. *Frontiers in Sports and Active Living*, 1, 70. <https://doi.org/10.3389/fspor.2019.00070>

Flockhart, M., & Larsen, F. J. (2024). Continuous glucose monitoring in endurance athletes: interpretation and relevance of measurements for improving performance and health. *Sports Medicine*, 54(2), 247–255. <https://doi.org/10.1007/s40279-023-01910-4>

Franch, J., Madsen, K., Djurhuus, M. S., & Pedersen, P. K. (1998). Improved running economy following intensified training correlates with reduced ventilatory demands. *Medicine and Science in Sports and Exercise*, 30(8), 1250–1256. <https://doi.org/10.1097/00005768-199808000-00011>

Furrer, R., Hawley, J. A., & Handschin, C. (2023). The molecular athlete: exercise physiology from mechanisms to medals. *Physiological Reviews*, 103(3), 1693–1787. <https://doi.org/10.1152/physrev.00017.2022>

Gibala M. (2009). Molecular responses to high-intensity interval exercise. *Applied Physiology, Nutrition, and Metabolism = Physiologie Appliquée, Nutrition et Métabolisme*, 34(3), 428–432. <https://doi.org/10.1139/H09-046>

Gojanovic, B., Shultz, R., Feihl, F., & Matheson, G. (2015). Overspeed HIIT in lower-body positive pressure treadmill improves running performance. *Medicine and Science in Sports and Exercise*, 47(12), 2571–2578. <https://doi.org/10.1249/MSS.0000000000000707>

Gonçalves, T. J. M., Boutilon, F., Lefebvre, S., Goffin, V., Iwatsubo, T., Wakabayashi, T., Oury, F., & Armand, A. S. (2019). Collagen XXV promotes myoblast fusion during myogenic differentiation and muscle formation. *Scientific Reports*, 9(1), 5878. <https://doi.org/10.1038/s41598-019-42296-6>

González-Mohíno, F., González-Ravé, J. M., Juárez, D., Fernández, F. A., Barragán Castellanos, R., & Newton, R. U. (2016). Effects of continuous and interval training on running economy, maximal aerobic speed and gait kinematics in recreational runners. *Journal of Strength and Conditioning Research*, 30(4), 1059–1066. <https://doi.org/10.1519/JSC.0000000000001174>

González-Mohíno, F., Santos-Concejero, J., Yustres, I., & González-Ravé, J. M. (2020). The effects of interval and continuous training on the oxygen cost of running in recreational runners: a systematic review and meta-analysis. *Sports Medicine*, 50(2), 283–294. <https://doi.org/10.1007/s40279-019-01201-x>

Grieb, A., Schmitt, A., Fragasso, A., Widmann, M., Mattioni Maturana, F., Burgstahler, C., Erz, G., Schellhorn, P., Nieß, A. M., & Munz, B. (2023). Skeletal muscle microRNA patterns in response to a single bout of exercise in females: biomarkers for subsequent training adaptation? *Biomolecules*, 13(6), 884. <https://doi.org/10.3390/biom13060884>

Guo, Y., Xiao, C., Zhao, K., He, Z., Liu, S., Wu, X., Shi, S., Chen, Z., & Shi, R. (2022). Physical exercise modalities for the management of heart failure with preserved ejection fraction: a systematic review and meta-analysis. *Journal of Cardiovascular Pharmacology*, 79(5), 698–710. <https://doi.org/10.1097/FJC.0000000000001254>

Hesketh, K., Shepherd, S. O., Strauss, J. A., Low, D. A., Cooper, R. J., Wagenmakers, A. J. M., & Cocks, M. (2019). Passive heat therapy in sedentary humans increases skeletal muscle capillarization and eNOS content but not mitochondrial density or GLUT4 content. *American Journal of Physiology. Heart and Circulatory Physiology*, 317(1), H114–H123. <https://doi.org/10.1152/ajpheart.00816.2018>

Higgins, J. P., & Green, S. (2011). *Cochrane handbook for systematic reviews of interventions version 5.1.0*. In The Cochrane Collaboration. Available at: www.handbook.cochrane.org; accessed on: 27 March 2024.

Higgins, J. P. T., Thomas, J., Chandler, J., Cumpston, M., Li, T., Page, M. J., Welch, V. A., & editors. (2023). *Cochrane Handbook for Systematic Reviews of Interventions Version 6.4*. 2023. Available at: www.training.cochrane.org/handbook; accessed on: 02 April 2024.

Higgins, J. P., Thompson, S. G., Deeks, J. J., & Altman, D. G. (2003). Measuring inconsistency in meta-analyses. *BMJ*, 327(7414), 557. <https://doi.org/10.1136/bmj.327.7414.557>

Holloszy, J. O., Rennie, M. J., Hickson, R. C., Conlee, R. K., & Hagberg, J. M. (1977). Physiological consequences of the biochemical adaptations to endurance exercise. *Annals of the New York Academy of Sciences*, 301, 440–450. <https://doi.org/10.1111/j.1749-6632.1977.tb38220.x>

Hoogeveen, A. R. (2000). The effect of endurance training on the ventilatory response to exercise in elite cyclists. *European Journal of Applied Physiology*, 82(2), 45–51. <https://doi.org/10.1007/s004210050650>

Hov, H., Wang, E., Lim, Y. R., Trane, G., Hemmingsen, M., Hoff, J., & Helgerud, J. (2023). Aerobic high-intensity intervals are superior to improve $\text{VO}_{2\text{max}}$ compared with sprint intervals in well-trained men. *Scandinavian Journal of Medicine and Science in Sports*, 33(2), 146–159. <https://doi.org/10.1111/sms.14251>

Jarstad, E., & Mamen, A. (2019). The performance and aerobic endurance effects of high-intensity versus moderate-intensity continuous running. *Applied Physiology, Nutrition, and Metabolism = Physiologie Appliquée, Nutrition et Métabolisme*, 44(9), 990–996. <https://doi.org/10.1139/apnm-2018-0575>

Jastrzębski, Z., Wakuluk-Lewandowska, D., Arslan, E., Kilit, B., Soylu, Y., & Radzimiński, Ł. (2025). Effects of eight-week game-based high-intensity interval training performed on different pitch dimensions on the level of physical capacity and time-motion responses in youth soccer players. *Journal of Human Kinetics*, 97, 157–168. <https://doi.org/10.5114/jhk/190842>

Jones, A. M., Kirby, B. S., Clark, I. E., Rice, H. M., Fulkerson, E., Wylie, L. J., Wilkerson, D. P., Vanhatalo, A., & Wilkins, B. W. (2021). Physiological demands of running at 2-hour marathon race pace. *Journal of Applied Physiology*, 130(2), 369–379. <https://doi.org/10.1152/japplphysiol.00647.2020>

Joyner, M. J., & Coyle, E. F. (2008). Endurance exercise performance: the physiology of champions. *Journal of Physiology*, 586(1), 35–44. <https://doi.org/10.1113/jphysiol.2007.143834>

Jung, M. E., Bourne, J. E., Beauchamp, M. R., Robinson, E., & Little, J. P. (2015). High-intensity interval training as an efficacious alternative to moderate-intensity continuous training for adults with prediabetes. *Journal of Diabetes Research*, 2015(1), 191595. <https://doi.org/10.1155/2015/191595>

Keir, D. A., Iannetta, D., Mattioni Maturana, F., Kowalchuk, J. M., & Murias, J. M. (2022). Identification of non-invasive exercise thresholds: methods, strategies, and an online App. *Sports Medicine*, 52(2), 237–255. <https://doi.org/10.1007/s40279-021-01581-z>

Khanna, G. L., & Manna, I. (2005). Supplementary effect of carbohydrate-electrolyte drink on sports performance, lactate removal & cardiovascular response of athletes. *Indian Journal of Medical Research*, 121(5), 665–669.

Kingsley, M. I., Wadsworth, D., Kilduff, L. P., McEneny, J., & Benton, D. (2005). Effects of phosphatidylserine on oxidative stress following intermittent running. *Medicine and Science in Sports and Exercise*, 37(8), 1300–1306. <https://doi.org/10.1249/01.mss.0000175306.05465.7e>

Kjøbsted, R., Hingst, J. R., Fentz, J., Foretz, M., Sanz, M. N., Pehmøller, C., Shum, M., Marette, A., Mounier, R., Treebak, J. T., Wojtaszewski, J. F. P., Viollet, B., & Lantier, L. (2018). AMPK in skeletal muscle function and metabolism. *FASEB Journal: Official Publication of the Federation of American Societies for Experimental Biology*, 32(4), 1741–1777. <https://doi.org/10.1096/fj.201700442R>

Lamboley, C. R., Rouffet, D. M., Dutka, T. L., McKenna, M. J., & Lamb, G. D. (2020). Effects of high-intensity intermittent exercise on the contractile properties of human type I and type II skeletal muscle fibers. *Journal of Applied Physiology*, 128(5), 1207–1216. <https://doi.org/10.1152/japplphysiol.00014.2020>

Laursen, P. B. (2010). Training for intense exercise performance: high-intensity or high-volume training? *Scandinavian Journal of Medicine & Science in Sports*, 20 Suppl 2, 1–10. <https://doi.org/10.1111/j.1600-0838.2010.01184.x>

Laursen, P. B., Marsh, S. A., Jenkins, D. G., & Coombes, J. S. (2007). Manipulating training intensity and volume in already well-trained rats: effect on skeletal muscle oxidative and glycolytic enzymes and buffering capacity. *Applied physiology, Nutrition, and Metabolism = Physiologie Appliquée, Nutrition et Métabolisme*, 32(3), 434–442. <https://doi.org/10.1139/H07-006>

Leite, C. D. F. C., Zovico, P. V. C., Rica, R. L., Barros, B. M., Machado, A. F., Evangelista, A. L., Leite, R. D., Barauna, V. G., Maia, A. F., & Bocalini, D. S. (2023). Exercise-induced muscle damage after a high-intensity interval exercise session: systematic review. *International Journal of Environmental Research and Public Health*, 20(22), 7082. <https://doi.org/10.3390/ijerph20227082>

Liang, W., Liu, C., Yan, X., Hou, Y., Yang, G., Dai, J., & Wang, S. (2024). The impact of sprint interval training versus moderate intensity continuous training on blood pressure and cardiorespiratory health in adults: a systematic review and meta-analysis. *PeerJ*, 12, e17064. <https://doi.org/10.7717/peerj.17064>

Litleskare, S., Enoksen, E., Sandvei, M., Støen, L., Stensrud, T., Johansen, E., & Jensen, J. (2020). Sprint interval running and continuous running produce training specific adaptations, despite a similar improvement of aerobic endurance capacity a randomized trial of healthy adults. *International Journal of Environmental Research and Public Health*, 17(11), 3865. <https://doi.org/10.3390/ijerph17113865>

Liu, B., Wu, J., Shi, Q., Hao, F., Xiao, W., Yu, J., Yu, F., & Ren, Z. (2022). Running economy and lower extremity stiffness in endurance runners: a systematic review and meta-analysis. *Frontiers in Physiology*, 13, 1059221. <https://doi.org/10.3389/fphys.2022.1059221>

Liu, H., Li, R., Zheng, W., Ramirez-Campillo, R., de Villarreal, E. S., & Zhang, M. (2024). The effect of combined strength, plyometric, and sprint training on repeated sprint ability in team-sport athletes: a systematic review and meta-analysis. *Journal of Sports Science & Medicine*, 23(4), 718–743. <https://doi.org/10.52082/jssm.2024.718>

Mandić, M., Hansson, B., Lovrić, A., Sundblad, P., Vollaard, N. B. J., Lundberg, T. R., Gustafsson, T., & Rullman, E. (2022). Improvements in maximal oxygen uptake after sprint-interval training coincide with increases in central hemodynamic factors. *Medicine and Science in Sports and Exercise*, 54(6), 944–952. <https://doi.org/10.1249/MSS.0000000000002872>

Massart, J., Sjögren, R. J. O., Egan, B., Garde, C., Lindgren, M., Gu, W., Ferreira, D. M. S., Katayama, M., Ruas, J. L., Barrès, R., O’Gorman, D. J., Zierath, J. R., & Krook, A. (2021). Endurance exercise training-responsive miR-19b-3p improves skeletal muscle glucose metabolism. *Nature Communications*, 12(1), 5948. <https://doi.org/10.1038/s41467-021-26095-0>

Maunder, E., Bradley, H. E., Deane, C. S., Hodgson, A. B., Jones, M., Joannis, S., Turner, A. M., Breen, L., Philp, A., & Wallis, G. A. (2021). Effects of short-term graded dietary carbohydrate intake on intramuscular and whole-body metabolism during moderate-intensity exercise. *Journal of Applied Physiology*, 131(1), 376–387. <https://doi.org/10.1152/japplphysiol.00811.2020>

Mayhew, T. P., Rothstein, J. M., Finucane, S. D., & Lamb, R. L. (1995). Muscular adaptation to concentric and eccentric exercise at equal power levels. *Medicine and Science in Sports and Exercise*, 27(6), 868–873. <https://doi.org/10.1249/00005768-199506000-00011>

Mølmen, K. S., Almquist, N. W., & Skattebo, Ø. (2025). Effects of exercise training on mitochondrial and capillary growth in human skeletal muscle: a systematic review and meta-regression. *Sports Medicine*, 55(1), 115–144. <https://doi.org/10.1007/s40279-024-02120-2>

Morais, J., Kilit, B., Arslan, E., Bragada, J., Soylu, Y., & Marinho, D. (2025). Effects of on-court tennis training combined with HIIT versus RST on aerobic capacity, speed, agility, jumping ability, and internal loads in young tennis players. *Journal of Human Kinetics*, 95, 173–185. <https://doi.org/10.5114/jhk/189691>

Morgan, D. W., Bransford, D. R., Costill, D. L., Daniels, J. T., Howley, E. T., & Krahenbuhl, G. S. (1995). Variation in the aerobic demand of running among trained and untrained subjects. *Medicine and Science in Sports and Exercise*, 27(3), 404–409.

Moritani, T., & deVries, H. A. (1979). Neural factors versus hypertrophy in the time course of muscle strength gain. *American Journal of Physical Medicine*, 58(3), 115–130.

Mougin, F., Simon-Rigaud, M. L., Davenne, D., Renaud, A., Garnier, A., Kantelip, J. P., & Magnin, P. (1991). Effects of sleep disturbances on subsequent physical performance. *European Journal of Applied Physiology and Occupational Physiology*, 63(2), 77–82. <https://doi.org/10.1007/BF00235173>

Munoz, I., Seiler, S., Alcocer, A., Carr, N., & Esteve-Lanao, J. (2015). Specific intensity for peaking: is race pace the best option? *Asian Journal of Sports Medicine*, 6(3), e24900. <https://doi.org/10.5812/asjsm.24900>

Nakagawa, S., & Cuthill, I. C. (2007). Effect size, confidence interval and statistical significance: a practical guide for biologists. *Biological Reviews of the Cambridge Philosophical Society*, 82(4), 591–605. <https://doi.org/10.1111/j.1469-185X.2007.00027.x>

Nibbeling, N., Daanen, H. A., Gerritsma, R. M., Hofland, R. M., & Oudejans, R. R. (2012). Effects of anxiety on running with and without an aiming task. *Journal of Sports Sciences*, 30(1), 11–19. <https://doi.org/10.1080/02640414.2011.617386>

Nuuttila, O. P., Nummela, A., Kyröläinen, H., Laukkanen, J., & Häkkinen, K. (2022). Physiological, perceptual, and performance responses to the 2-week block of high- versus low-intensity endurance training. *Medicine and Science in Sports and Exercise*, 54(5), 851–860. <https://doi.org/10.1249/MSS.0000000000002861>

Page, M. J., McKenzie, J. E., Bossuyt, P. M., & et al. (2021). The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ*, 372, n71. <https://doi.org/10.1136/bmj.n71>

Parmar, A., Jones, T. W., & Hayes, P. R. (2021). The dose-response relationship between interval-training and $\text{VO}_{2\text{max}}$ in well-trained endurance runners: a systematic review. *Journal of Sports Sciences*, 39(12), 1410–1427. <https://doi.org/10.1080/02640414.2021.1876313>

Peake, J. M., Tan, S. J., Markworth, J. F., Broadbent, J. A., Skinner, T. L., & Cameron-Smith, D. (2014). Metabolic and hormonal responses to isoenergetic high-intensity interval exercise and continuous moderate-intensity exercise. *American Journal of Physiology. Endocrinology and Metabolism*, 307(7), E539–E552. <https://doi.org/10.1152/ajpendo.00276.2014>

Perry, C. G., Heigenhauser, G. J., Bonen, A., & Spriet, L. L. (2008). High-intensity aerobic interval training increases fat and carbohydrate metabolic capacities in human skeletal muscle. *Applied Physiology, Nutrition, and Metabolism = Physiologie Appliquée, Nutrition et Métabolisme*, 33(6), 1112–1123. <https://doi.org/10.1139/H08-097>

Phillips, S. M., & Van Loon, L. J. C. (2011). Dietary protein for athletes: from requirements to metabolic advantage. *Applied Physiology, Nutrition, and Metabolism*, 36(5), 683–694.

Rago, V., Krustrup, P., & Mohr, M. (2022). Performance and submaximal adaptations to additional speed-endurance training Vs. continuous moderate-intensity aerobic training in male endurance athletes. *Journal of Human Kinetics*, 83, 277–285. <https://doi.org/10.2478/hukin-2022-0060>

Rakobowchuk, M., Tanguay, S., Burgomaster, K. A., Howarth, K. R., Gibala, M. J., & MacDonald, M. J. (2008). Sprint interval and traditional endurance training induce similar improvements in peripheral arterial stiffness and flow-mediated dilation in healthy humans. *American Journal of Physiology. Regulatory Integrative and Comparative Physiology*, 295(1), R236–R242. <https://doi.org/10.1152/ajpregu.00069.2008>

Ramos, J. S., Dalleck, L. C., Tjonna, A. E., Beetham, K. S., & Coombes, J. S. (2015). The impact of high-intensity interval training versus moderate-intensity continuous training on vascular function: a systematic review and meta-analysis. *Sports Medicine*, 45(5), 679–692. <https://doi.org/10.1007/s40279-015-0321-z>

Reuter, M., Rosenberger, F., Barz, A., Venhorst, A., Blanz, L., Roecker, K., & Meyer, T. (2024). Effects on cardiorespiratory fitness of moderate-intensity training vs. energy-matched training with increasing intensity. *Frontiers in Sports and Active Living*, 5, 1298877. <https://doi.org/10.3389/fspor.2023.1298877>

Rimmele, U., Seiler, R., Marti, B., Wirtz, P. H., Ehlert, U., & Heinrichs, M. (2009). The level of physical activity affects adrenal and cardiovascular reactivity to psychosocial stress. *Psychoneuroendocrinology*, 34(2), 190–198. <https://doi.org/10.1016/j.psyneuen.2008.08.023>

Rognmo, Ø., Moholdt, T., Bakken, H., Hole, T., Mølstad, P., Myhr, N. E., Grimsmo, J., & Wisløff, U. (2012). Cardiovascular risk of high- versus moderate-intensity aerobic exercise in coronary heart disease patients. *Circulation*, 126(12), 1436–1440. <https://doi.org/10.1161/circulationaha.112.123117>

Roy, M., Williams, S. M., Brown, R. C., Meredith-Jones, K. A., Osborne, H., Jospe, M., & Taylor, R. W. (2018). High-intensity interval training in the real world: outcomes from a 12-month intervention in overweight adults. *Medicine and Science in Sports and Exercise*, 50(9), 1818–1826. <https://doi.org/10.1249/MSS.0000000000001642>

Ruegsegger, G. N., Pataky, M. W., Simha, S., Robinson, M. M., Klaus, K. A., & Nair, K. S. (2023). High-intensity aerobic, but not resistance or combined, exercise training improves both cardiometabolic health and skeletal muscle mitochondrial dynamics. *Journal of Applied Physiology*, 135(4), 763–774. <https://doi.org/10.1152/japplphysiol.00405.2023>

Santos, A., Braaten, K., MacPherson, M., Vasconcellos, D., Vis-Dunbar, M., Lonsdale, C., Lubans, D., & Jung, M. E. (2023). Rates of compliance and adherence to high-intensity interval training: a systematic review and meta-analyses. *International Journal of Behavioral Nutrition and Physical Activity*, 20(1), 134. <https://doi.org/10.1186/s12966-023-01535-w>

Saunders, P. U., Pyne, D. B., Telford, R. D., & Hawley, J. A. (2004). Factors affecting running economy in trained distance runners. *Sports Medicine*, 34(7), 465–485. <https://doi.org/10.2165/00007256-200434070-00005>

Sawyer, B. J., Tucker, W. J., Bhammar, D. M., Ryder, J. R., Sweazea, K. L., & SGAesser, G. A. (2016). Effects of high-intensity interval training and moderate-intensity continuous training on endothelial function and cardiometabolic risk markers in obese adults. *Journal of Applied Physiology*, 121(1), 279–288. <https://doi.org/10.1152/japplphysiol.00024.2016>

Schaun, G. Z., Pinto, S. S., Silva, M. R., Dolinski, D. B., & Alberton, C. L. (2018). Whole-body high-intensity interval training induce similar cardiorespiratory adaptations compared with traditional high-intensity interval training and moderate-intensity continuous training in healthy men. *Journal of Strength and Conditioning Research*, 32(10), 2730–2742. <https://doi.org/10.1519/JSC.0000000000002594>

Schommer, N. C., Hellhammer, D. H., & Kirschbaum, C. (2003). Dissociation between reactivity of the hypothalamus-pituitary-adrenal axis and the sympathetic-adrenal-medullary system to repeated psychosocial stress. *Psychosomatic Medicine*, 65(3), 450–460. <https://doi.org/10.1097/01.psy.0000035721.12441.17>

Seiler, K. S., & Kjerland, G. Ø. (2006). Quantifying training intensity distribution in elite endurance athletes: is there evidence for an optimal distribution? *Scandinavian Journal of Medicine & Science in Sports*, 16(1), 49–56. <https://doi.org/10.1111/j.1600-0838.2004.00418.x>

Seiler, S., Haugen, O., & Kuffel, E. (2007). Autonomic recovery after exercise in trained athletes: intensity and duration effects. *Medicine and Science in Sports and Exercise*, 39(8), 1366–1373. <https://doi.org/10.1249/mss.0b013e318060f17d>

Seo, M., Park, T., & Jung, H. C. (2024). Sex differences in heart rate variability and vascular function following high-Intensity interval training in young adults. *Journal of Human Kinetics*, 90, 89–100. <https://doi.org/10.5114/jhk/170964>

Shaw, A. J., Ingham, S. A., Fudge, B. W., & Folland, J. P. (2013). The reliability of running economy expressed as oxygen cost and energy cost in trained distance runners. *Applied Physiology, Nutrition, and Metabolism*, 38(12), 1268–72. <https://doi.org/10.1139/apnm-2013-0055>

Silva, R., Damasceno, M., Cruz, R., Silva-Cavalcante, M. D., Lima-Silva, A. E., Bishop, D. J., & Bertuzzi, R. (2017). Effects of a 4-week high-intensity interval training on pacing during 5-km running trial. *Brazilian Journal of Medical and Biological Research*, 50(12), e6335. <https://doi.org/10.1590/1414-431X20176335>

Skelly, L. E., Gillen, J. B., Frankish, B. P., MacInnis, M. J., Godkin, F. E., Tarnopolsky, M. A., Murphy, R. M., & Gibala, M. J. (2021). Human skeletal muscle fiber type-specific responses to sprint interval and moderate-intensity continuous exercise: acute and training-induced changes. *Journal of Applied Physiology*, 130(4), 1001–1014. <https://doi.org/10.1152/japplphysiol.00862.2020>

Skovgaard, C., Almquist, N.W., & Bangsbo, J. (2018a). The effect of repeated periods of speed endurance training on performance, running economy, and muscle adaptations. *Scandinavian Journal of Medicine & Science in Sports*, 28(2), 381–390. <https://doi.org/10.1111/sms.12916>

Skovgaard, C., Christiansen, D., Christensen, P. M., Almquist, N. W., Thomassen, M., & Bangsbo, J. (2018b). Effect of speed endurance training and reduced training volume on running economy and single muscle fiber adaptations in trained runners. *Physiological Reports*, 6(3), e13601. <https://doi.org/10.14814/phy2.13601>

Sperlich, B., Zinner, C., Heilemann, I., Kjendlie, P. L., Holmberg, H. C., & Mester, J. (2010). High-intensity interval training improves VO_{2peak}, maximal lactate accumulation, time trial and competition performance in 9–11-year-old swimmers. *European Journal of Applied Physiology*, 110(5), 1029–1036. <https://doi.org/10.1007/s00421-010-1586-4>

Stögg, T. L., & Sperlich, B. (2015). The training intensity distribution among well-trained and elite endurance athletes. *Frontiers in Physiology*, 6, 295. <https://doi.org/10.3389/fphys.2015.00295>

Tanji, F., & Nabekura, Y. (2019). Oxygen uptake and respiratory exchange ratio relative to the lactate threshold running in well-trained distance runners. *Journal of Sports Medicine and Physical Fitness*, 59(6), 895–901. <https://doi.org/10.23736/S0022-4707.18.08828-X>

Taylor, J. L., Holland, D. J., Keating, S. E., Leveritt, M. D., Gomersall, S. R., Rowlands, A. V., Bailey, T. G., & Coombes, J. S. (2020). Short-term and long-term feasibility, safety, and efficacy of high-intensity interval training in cardiac rehabilitation: the Fit heart study randomized clinical trial. *JAMA Cardiology*, 5(12), 1382–1389. <https://doi.org/10.1001/jamacardio.2020.3511>

Thomas, C., Sirvent, P., Perrey, S., Raynaud, E., & Mercier, J. (2004). Relationships between maximal muscle oxidative capacity and blood lactate removal after supramaximal exercise and fatigue indexes in humans. *Journal of Applied Physiology*, 97(6), 2132–2138. <https://doi.org/10.1152/japplphysiol.00387.2004>

Torma, F., Gombos, Z., Jokai, M., Takeda, M., Mimura, T., & Radak, Z. (2019). High intensity interval training and molecular adaptive response of skeletal muscle. *Sports Medicine and Health Science*, 1(1), 24–32. <https://doi.org/10.1016/j.smhs.2019.08.003>

Unhjem R. J. (2024). Changes in running economy and attainable maximal oxygen consumption in response to prolonged running: the impact of training status. *Scandinavian Journal of Medicine & Science in Sports*, 34(5), e14637. <https://doi.org/10.1111/sms.14637>

Van Hooren, B., Jukic, I., Cox, M., Frenken, KG., Bautista, I., & Moore, IS. (2024). The relationship between running biomechanics and running economy: a systematic review and meta-analysis of observational studies. *Sports Medicine*, 54(5), 1269–1316. <https://doi.org/10.1007/s40279-024-01997-3>

Vitale, K. C., Owens, R., Hopkins, S. R., & Malhotra, A. (2019). Sleep hygiene for optimizing recovery in athletes: review and recommendations. *International Journal of Sports Medicine*, 40(8), 535–543. <https://doi.org/10.1055/a-0905-3103>

Vuorimaa, T., Vasankari, T., & Rusko, H. (2000). Comparison of physiological strain and muscular performance of athletes during two intermittent running exercises at the velocity associated with $VO_{2\text{max}}$. *International Journal of Sports Medicine*, 21(2), 96–101. <https://doi.org/10.1055/s-2000-8867>

Wang, F., Wang, X., Liu, Y., & Zhang, Z. (2021). Effects of exercise-induced ROS on the pathophysiological functions of skeletal muscle. *Oxidative Medicine and Cellular Longevity*, 2021, 3846122. <https://doi.org/10.1155/2021/3846122>

Wang, T. Y., Ho, C. F., Chan, K. H., Lee, W. C., & Hsu, M. C. (2012). Effects of consecutive 7-day high- versus moderate-intensity training on endurance determinants and muscle damage in basketball players. *International SportMed Journal*, 13(1), 18–28.

Wang, Z., & Wang, J. (2024). The effects of high-intensity interval training versus moderate-intensity continuous training on athlete aerobic endurance performance parameters. *European Journal of Applied Physiology*, 124(8), 2235–2249. <https://doi.org/10.1007/s00421-024-05532-0>

Webb, H. E., Weldy, M. L., Fabianke-Kadue, E. C., Orndorff, G. R., Kamimori, G. H., & Acevedo, E. O. (2008). Psychological stress during exercise: cardiorespiratory and hormonal responses. *European Journal of Applied Physiology*, 104(6), 973–981. <https://doi.org/10.1007/s00421-008-0852-1>

Wen, D., Utesch, T., Wu, J., Robertson, S., Liu, J., Hu, G., & Chen, H. (2019). Effects of different protocols of high intensity interval training for $VO_{2\text{max}}$ improvements in adults: a meta-analysis of randomised controlled trials. *Journal of Science and Medicine in Sport*, 22(8), 941–947. <https://doi.org/10.1016/j.jsams.2019.01.013>

Weston, A. R., Myburgh, K. H., Lindsay, F. H., Dennis, S. C., Noakes, T. D., & Hawley, J. A. (1997). Skeletal muscle buffering capacity and endurance performance after high-intensity interval training by well-trained cyclists. *European Journal of Applied Physiology and Occupational Physiology*, 75(1), 7–13. <https://doi.org/10.1007/s004210050119>

Widmann, M., Mattioni Maturana, F., Burgstahler, C., Erz, G., Schellhorn, P., Fragasso, A., Schmitt, A., Nieß, A. M., & Munz, B. (2022). MiRNAs as markers for the development of individualized training regimens: a pilot study. *Physiological Reports*, 10(5), e15217. <https://doi.org/10.14814/phy2.15217>

Wiewelhove, T., Fernandez-Fernandez, J., Raeder, C., Kappenstein, J., Meyer, T., Kellmann, M., Pfeiffer, M., & Ferrauti, A. (2016). Acute responses and muscle damage in different high-intensity interval running protocols. *Journal of Sports Medicine and Physical Fitness*, 56(5), 606–615.

Yoshida, T., Udo, M., Iwai, K., & Yamaguchi, T. (1993). Physiological characteristics related to endurance running performance in female distance runners. *Journal of Sports Sciences*, 11(1), 57–62. <https://doi.org/10.1080/02640419308729964>

Zinner, C., Schäfer Olstad, D., & Sperlich, B. (2018). Mesocycles with different training intensity distribution in recreational runners. *Medicine and Science in Sports and Exercise*, 50(8), 1641–1648. <https://doi.org/10.1249/MSS.0000000000001599>